

PERMUTATIONS IN TWO DIMENSIONS THAT MAXIMALLY SEPARATE NEIGHBORS

MOHAMMED ALBOW, JEFF EDGINGTON, MARIO LOPEZ, AND PETR VOJTĚCHOVSKÝ

ABSTRACT. We characterize all permutations on even-by-even grids that maximally separate neighboring vertices. More precisely, let n_1, n_2 be positive even integers, let $I(n_1, n_2) = \{1, \dots, n_1\} \times \{1, \dots, n_2\}$ be the $n_1 \times n_2$ grid, let d be the L_1 metric on $I(n_1, n_2)$, and let $N = \{\{x, y\} \in I(n_1, n_2) \times I(n_1, n_2) : d(x, y) = 1\}$ be the set of neighbors in $I(n_1, n_2)$. We characterize all permutations π of $I(n_1, n_2)$ that maximize $\sum_{\{x, y\} \in N} d(\pi(x), \pi(y))$.

1. INTRODUCTION

The *total displacement* $\sum_{i=1}^n |\pi(i) - i|$ of a permutation π on $\{1, 2, \dots, n\}$ was investigated by Diaconis and Graham [5], Gallero et al [6], Daly and Vojtěchovský [4], Guay-Paquet and Petersen [7], Bärtschi et al [2], and others. The concept appears in the literature under various names and with subtle variations; the above definition and terminology is due to Knuth [9, Problem 5.1.1.28]. The displacements $|\pi(i) - i|$ play a role in the construction of interleavers for turbo codes [3].

In this paper we study a quantity similar to total displacement that measures how much permutations *separate* neighboring points. Let X be a finite set, d a metric on X , and $N = \{\{x, y\} \in X \times X : d(x, y) = 1\}$ the set of (unit) neighbors in X . For a permutation π of X , the quantity

$$\frac{1}{|N|} \sum_{\{x, y\} \in N} d(\pi(x), \pi(y))$$

yields the average distance between π -images of neighboring points. Without the normalizing factor $1/|N|$, we obtain the function

$$(1.1) \quad f : S_{(X, d)} \rightarrow \mathbb{N}, \quad f(\pi) = \sum_{\{x, y\} \in N} d(\pi(x), \pi(y)).$$

We are interested in the properties of f , in particular in the following questions: *What is the maximal value of f ? For which permutations π of X is the maximum of f attained?*

These and other questions were studied and answered in [4] for the simplest case of a one-dimensional interval $X = \{1, 2, \dots, n\}$ equipped with the L_1 metric $d(i, j) = |i - j|$, cf. [4, Theorem 3.1] or Theorem 4.4. Here we answer the two questions for permutations on even-by-even grids $I(2t_1, 2t_2)$ equipped with the two-dimensional L_1 metric. The solution is much more intricate than in one dimension and can be roughly described as follows:

2010 *Mathematics Subject Classification.* 05A05.

Key words and phrases. Permutation in two dimensions, neighbors in graphs, separation of neighboring vertices, total displacement of a permutation.

Research partially supported by the Simons Foundation Collaboration Grant 210176 to Petr Vojtěchovský.

- the four entries at the center of the grid must be moved into the four corners,
- further concentric layers surrounding the center must be moved onto the boundary of the grid,
- images must maintain a double checkerboard pattern as much as possible, in the sense that if the images $x = (x_1, x_2)$, $y = (y_1, y_2)$ are neighbors, then the values x_1 , x_2 must not be both small (in $[1, \dots, t_1]$) nor both large, and similarly for y_1 , y_2 ,
- the unavoidable defects in the double checkerboard pattern must be located along the line that partitions the grid into two equal halves along the shorter dimension,
- the values involved in the defects must alternate between t_1 and $t_1 + 1$, or between t_2 and $t_2 + 1$.

See Construction 3.1 for a more precise description of the solutions, Figure 2 for an example on the 6×6 grid, Theorem 3.7 for a complete description of the solutions in the generic case, and the comments following Theorem 3.7 for the four exceptional cases.

The paper concludes with comments on the general two-dimensional case and on higher-dimensional cases, where the solutions are suspected to behave quite differently.

A playful example that motivates the two-dimensional problem is as follows. *Students are about to take a test. Assuming that the students occupy all classroom seats arranged in a rectangular grid, how can the students be reseated so that the average distance to their previous neighbors is maximized?* Every experienced instructor will immediately recognize the practical importance of this problem.

We are not aware of an engineering application of our result. Nevertheless, two-dimensional interleavers have been considered for the transmission of two-dimensional data in noisy channels [8], as were two-dimensional constructions of regular interleavers for turbo codes [3].

1.1. Notation. All sets in this paper are treated as multisets, they will be denoted by uppercase letters, and their cardinalities will be denoted by the corresponding lowercase letters. For instance, a is the cardinality of A . We write $\sum A$ as a shorthand for $\sum_{x \in A} x$, and $k*x$ to indicate that the element x appears k times. For instance, $\{3*2, 2*4\} = \{2, 2, 2, 4, 4\}$.

We denote the coordinates of $x = (i, j)$ of $\mathbb{Z} \times \mathbb{Z}$ by $x_1 = i$ and $x_2 = j$. The subscript 1 refers to rows, while the subscript 2 refers to columns. If A is a submultiset of $\mathbb{Z} \times \mathbb{Z}$ and $i \in \{1, 2\}$, we implicitly define

$$A_i = \{x_i : x \in A\}.$$

Moreover, given $A_1, A_2 \subseteq \mathbb{Z}$, we implicitly define

$$A_* = A_1 \cup A_2.$$

Note that $A \subseteq \mathbb{Z} \times \mathbb{Z}$, while $A_* \subseteq \mathbb{Z}$.

Fix positive integers n_1, n_2 . Let $I = I(n_1, n_2) = \{1, \dots, n_1\} \times \{1, \dots, n_2\}$ be the $n_1 \times n_2$ grid. We will visualize a permutation π of I as an $n_1 \times n_2$ array with the coordinate $(1, 1)$ in the top left corner, where we place (k, ℓ) in row i and column j if and only if $\pi(i, j) = (k, \ell)$.

Given a permutation π of I , let

$$C = \{\pi(1, 1), \pi(1, n_2), \pi(n_1, 1), \pi(n_1, n_2)\}$$

be the values of π on the four corners of I , and let

$$B = \{\pi(i, j) : i \in \{1, n_1\} \text{ or } j \in \{1, n_2\}\} \setminus C$$

be the values of π on the boundary of I excluding the four corners.

Consider the L_1 metric d on I defined by

$$d((i, j), (k, \ell)) = |i - k| + |j - \ell|.$$

Note that the corners have two neighbors, other boundary cells have three neighbors, and interior cells have four neighbors in I .

As a special case of the general definition (1.1), for a permutation π of I let

$$(1.2) \quad \begin{aligned} f(\pi) &= \sum_{\substack{1 \leq i < n_1 \\ 1 \leq j \leq n_2}} (|\pi(i, j)_1 - \pi(i+1, j)_1| + |\pi(i, j)_2 - \pi(i+1, j)_2|) \\ &+ \sum_{\substack{1 \leq i \leq n_1 \\ 1 \leq j < n_2}} (|\pi(i, j)_1 - \pi(i, j+1)_1| + |\pi(i, j)_2 - \pi(i, j+1)_2|). \end{aligned}$$

Let $S = S(n_1, n_2) = \{4 * (i, j) : (i, j) \in I(n_1, n_2)\}$ and recall that $S_1 = \{4n_2 * i : 1 \leq i \leq n_1\}$, $S_2 = \{4n_1 * i : 1 \leq i \leq n_2\}$ and S_* are implicitly defined.

Finally, the entries $\pi(i, j)_k$ of the combined sums of (1.2) form a submultiset of S_* . Therefore, there are suitable submultisets $S_i^+ = S_i^+(\pi)$, $S_i^- = S_i^-(\pi)$ of S_i such that

$$(1.3) \quad f(\pi) = \sum S_1^+ - \sum S_1^- + \sum S_2^+ - \sum S_2^-.$$

Note that $s_i^+ = s_i^- = n_1(n_2 - 1) + n_2(n_1 - 1)$.

Here is a small example:

Example 1.1. Consider the permutation π of $I(2, 3)$ given by

(2, 2)	(1, 3)	(2, 1)
(1, 2)	(1, 1)	(2, 3)

Then

$$\begin{aligned} f(\pi) &= (|2 - 1| + |2 - 3|) + (|1 - 2| + |3 - 1|) + (|1 - 1| + |2 - 1|) + (|1 - 2| + |1 - 3|) \\ &+ (|2 - 1| + |2 - 2|) + (|1 - 1| + |3 - 1|) + (|2 - 2| + |1 - 3|) = 14. \end{aligned}$$

Upon separating the row and column coordinates, we get

$$\begin{aligned} f(\pi) &= (|2 - 1| + |1 - 2| + |1 - 1| + |1 - 2| + |2 - 1| + |1 - 1| + |2 - 2|) \\ &+ (|2 - 3| + |3 - 1| + |2 - 1| + |1 - 3| + |2 - 2| + |3 - 1| + |1 - 3|). \end{aligned}$$

Finally, upon simplifying the absolute values and collecting positive and negative entries, we obtain

$$\begin{aligned} f(\pi) &= (2 + 2 + 1 + 2 + 2 + 1 + 2) - (1 + 1 + 1 + 1 + 1 + 1 + 2) \\ &+ (3 + 3 + 2 + 3 + 2 + 3 + 3) - (2 + 1 + 1 + 1 + 2 + 1 + 1) \\ &= \sum S_1^+ - \sum S_1^- + \sum S_2^+ - \sum S_2^-, \end{aligned}$$

where $S_1^+ = \{2 * 1, 5 * 2\}$, $S_1^- = \{6 * 1, 1 * 2\}$, $S_2^+ = \{2 * 2, 5 * 3\}$ and $S_2^- = \{5 * 1, 2 * 2\}$ are the desired multisets.

1.2. **An unattainable upper bound.** From now on suppose that $n_1 = 2t_1$ and $n_2 = 2t_2$. For $A_i \subseteq S_i$, let

$$A_i^< = \{u \in A_i : 1 \leq u \leq t_i\}, \quad A_i^> = \{u \in A_i : t_i < u \leq n_i\}.$$

We will refer to the elements of $A_i^<$ as the *small* values of A_i , and to the elements of $A_i^>$ as the *large* values of A_i . We have $A_i^< \cup A_i^> = A_i$, of course.

As a special case we obtain the multisets $S_i^<$, $S_i^>$ with cardinalities $s_i^< = s_i^> = 2n_1n_2$. Whenever two subsets $A_i^< \subseteq S_i^<$ and $A_i^> \subseteq S_i^>$ are given, we will assume that $A_i = A_i^< \cup A_i^>$ is implicitly defined.

Lemma 1.2. *Let π be a permutation of $I(n_1, n_2)$ with $n_i = 2t_i$. Then*

$$f(\pi) < \sum S_1^> - \sum S_1^< + \sum S_2^> - \sum S_2^<.$$

Proof. The equality (1.3) holds for suitable submultisets S_i^+ , S_i^- of S_i . We have $s_i^+ = s_i^- = n_1(n_2 - 1) + n_2(n_1 - 1) < 2n_1n_2 = s_i^> = s_i^<$. Let X_i^+ be the multiset consisting of the s_i^+ largest values in $S_i^>$, and X_i^- the multiset consisting of the s_i^- smallest values in $S_i^<$. Then, for $i \in \{1, 2\}$, we have

$$\sum S_i^> - \sum S_i^< > \sum X_i^+ - \sum X_i^- \geq \sum S_i^+ - \sum S_i^-.$$

Summing up the inequalities for $i = \{1, 2\}$ finishes the proof. \square

The upper bound of Lemma 1.2 cannot be achieved for two reasons. First, due to the boundary of the grid, not every symbol from S_i is used in $f(\pi)$ with the maximal multiplicity. Second, to count all large values with positive signs and all small values with negative signs in (1.3), the permutation π would have to form a checkerboard with respect to the sets $S_1^<$, $S_1^>$ in the first coordinate, and another checkerboard with respect to the sets $S_2^<$, $S_2^>$ in the second coordinate. This is not possible. Indeed, once we place an entry from $S_1^> \times S_2^>$ in the array defining π , we must continue the double checkerboard pattern by alternating entries from $S_1^> \times S_2^>$ with entries from $S_1^< \times S_2^<$. Since an entry from $S_1^< \times S_2^>$, say, has to be used eventually, it will be adjacent to either an entry from $S_1^> \times S_2^>$ or to an entry from $S_1^< \times S_2^<$, and it will break the pattern in one of the coordinates.

2. CORNERS, BOUNDARIES AND DEFECTS IN THE DOUBLE CHECKERBOARD PATTERN

2.1. **Defects.** To investigate the flaws in the double checkerboard pattern, we define the *defect* multisets

$$D_i^< = \{\max\{\pi(x)_i, \pi(y)_i\} : \{x, y\} \in N, \{\pi(x)_i, \pi(y)_i\} \subseteq S_i^<\},$$

$$D_i^> = \{\min\{\pi(x)_i, \pi(y)_i\} : \{x, y\} \in N, \{\pi(x)_i, \pi(y)_i\} \subseteq S_i^>\},$$

for $i \in \{1, 2\}$.

The definition of $D_1^<$ is justified as follows. If x, y are neighboring cells and if the two values $u = \pi(x)_1$, $v = \pi(y)_1$ are both small, say with $u < v$, then these two values will contribute to $f(\pi)$ by $-u + v$, while in the ideal (and globally unattainable) situation we would like them to contribute by $-u - v$. We therefore keep track of the value with the incorrect sign, namely $v = \max\{u, v\}$. Similarly for the sets $D_1^>$, $D_2^<$, $D_2^>$.

From now on, we will color cells with values in $S_1^> \times S_2^>$ dark blue, $S_1^< \times S_2^<$ light blue, $S_1^> \times S_2^<$ dark red, and $S_1^< \times S_2^>$ light red. We will call the (blue) values in $(S_1^> \times S_2^>) \cup (S_1^< \times S_2^<)$ *homogeneous*, and the (red) values in $(S_1^> \times S_2^<) \cup (S_1^< \times S_2^>)$ *heterogeneous*.

Let us give another example to elucidate many of the concepts defined so far.

Example 2.1. Consider the permutation π of $I(4, 6)$ defined by

(2, 6)	(3, 1)	(1, 1)	(3, 4)	(1, 4)	(3, 6)
(4, 3)	(2, 5)	(3, 2)	(1, 2)	(3, 5)	(2, 2)
(2, 4)	(4, 2)	(1, 3)	(4, 6)	(2, 1)	(4, 4)
(4, 1)	(1, 6)	(3, 3)	(1, 5)	(4, 5)	(2, 3)

Since the values inside the homogeneous (resp. heterogeneous) region happen to form a double checkerboard in this example, the defects can occur only where the two regions meet.

A quick inspection shows that we have $D_1 = \emptyset$, $D_2^< = D_*^< = \{1, 2, 2, 3, 3, 3\}$, $D_2^> = D_*^> = \{4, 4, 4, 5, 5\}$, $D_2 = D_* = \{1, 2 * 2, 3 * 3, 3 * 4, 2 * 5\}$, $C_1^< = \{2, 2\}$, $C_1^> = \{3, 4\}$, and so on.

The following, rather obvious lemma will be used to minimize the number of defects on the grid.

Lemma 2.2. *Let $n_1 \leq n_2$ be positive even integers. Consider $I(n_1, n_2)$ as a graph where two vertices x, y are joined by an edge if and only if $\{x, y\} \in N$. Color a half of the vertices red and the other half blue. Then there are at least n_1 red-blue edges.*

Suppose further that there are precisely n_1 red-blue edges. If $n_1 < n_2$ then the grid can be evenly split by a vertical cut into blue and red connected components. If $n_1 = n_2$ then the grid can be evenly split by a vertical cut or a horizontal cut into blue and red connected components.

Proof. Let us first show that the number e of red-blue edges satisfies $e \geq n_1$. If no row is monochromatic then $e \geq n_1$. If no column is monochromatic then $e \geq n_2 \geq n_1$. We can therefore assume that there exists a monochromatic row and a monochromatic column, without loss of generality forming a red cross. Let r (resp. c) be the number of rows (resp. columns) containing a blue vertex. Any row or column containing a blue vertex intersects the red cross, hence $r + c \leq e$. We also have $rc \geq n_1 n_2 / 2$, the number of blue vertices. Therefore, $n_1^2 \leq n_1 n_2 \leq 2rc < (r + c)^2 \leq e^2$, and $e > n_1$ follows.

Suppose that $e = n_1$. If no row is monochromatic then every column is monochromatic (else $e > n_1$) and we can see easily that the grid can be evenly split by a vertical cut into two monochromatic connected components. If there is a monochromatic row, then no column is monochromatic (else $e > n_1$ as in the first paragraph), which forces $n_1 = n_2$, and we conclude the proof using a transposed argument. \square

Corollary 2.3. *Every permutation of $I(2t_1, 2t_2)$ contains at least $2t_1$ defects, i.e., $d_* \geq 2t_1$.*

Proof. Color (i, j) blue if $\pi(i, j)$ is homogeneous, and red otherwise. By Lemma 2.2 there are at least $2t_1$ red-blue edges, each contributing a defect. For instance, an edge between $S_1^< \times S_2^<$ and $S_1^> \times S_2^<$ contributes a column defect. \square

2.2. An exact formula. The value of $f(\pi)$ can be expressed exactly in terms of the corners, boundaries and defects. To that effect, let

$$(2.1) \quad g(\pi) = \left(\sum B_*^> + 2 \sum C_*^> + 2 \sum D_*^> \right) - \left(\sum B_*^< + 2 \sum C_*^< + 2 \sum D_*^< \right).$$

Proposition 2.4. *Let π be a permutation of $I(2t_1, 2t_2)$. Then*

$$f(\pi) = \left(\sum S_*^> - \sum S_*^< \right) - g(\pi).$$

Proof. Consider a cell x with $\pi(x) = (i, j)$ and $j \in S_2^>$. The value j is counted $8t_1$ times in $\sum S_2^>$. We must make the following modifications to $f(\pi)$ on account of x . If $x \in B$ then j must be subtracted once from $\sum S_2^>$ because x has only three neighbors. If $x \in C$ then j must be subtracted twice from $\sum S_2^>$ because x has only two neighbors. If x participates in a defect in the second coordinate (as illustrated by the neighboring values $(1, 4)$ and $(3, 5)$ in Example 2.1), there is some neighbor y of x such that $\pi(y) = (i', j')$ with $j' \in S_2^>$. This defect contributes $k = \min\{j, j'\}$ to $D_2^>$, and we must subtract k two times from $\sum S_2^>$ because of the sign change in the defect.

Similarly for $j \in S_2^<$, except that the value has to be added to $-\sum S_2^<$. Analogously for the first coordinate. \square

Maximizing $f(\pi)$ is thus equivalent to minimizing the positive quantity $g(\pi)$. The following result shows that the number of positive summands in (2.1) is equal to the number of negative summands in (2.1).

Proposition 2.5. *Let π be a permutation of $I(2t_1, 2t_2)$. Then*

$$b_i^> + 2c_i^> + 2d_i^> = b_i^< + 2c_i^< + 2d_i^<$$

for $i \in \{1, 2\}$.

Proof. Throughout the proof, if we deal with two permutations π and π' , then $b_i^>$ and similar quantities will refer to π , while $b_i'^>$ and similar quantities will refer to π' .

We will prove the result for $i = 1$. The case $i = 2$ is similar. Place the grid on a torus by identifying the opposite sides, and interpret the result as having no corners and boundaries, i.e., $c_* = b_* = 0$. Fix any initial permutation on the torus for which small and large values alternate in the first coordinate, that is, $d_1 = 0$. Such a permutation exists since both dimensions are even.

Any other permutation on the torus is obtained from the initial permutation by a sequence of successive swaps of two cells. Suppose that the result holds for π and let π' be the permutation obtained from π by swapping yet another pair of cells x, y . If $\pi(x)_1, \pi(y)_1$ are both small or both large, then $d_1'^< = d_1^<$ and $d_1'^> = d_1^>$. Otherwise let us assume without loss of generality that $(\pi(x)_1, \pi(y)_1) \in S_1^< \times S_1^>$. Let $z_x^<$ (resp. $z_x^>$) denote the number of neighbors of x in the grid of π that have a small entry (resp. a large entry) in the first coordinate, and define $z_y^<, z_y^>$ similarly. Then $d_1'^< = d_1^< - z_x^< + z_y^<$ and $d_1'^> = d_1^> - z_y^> + z_x^>$. Thus $d_1'^< - d_1'^> = d_1^< - d_1^> - z_x^< - z_x^> + z_y^< + z_y^> = d_1^< - d_1^>$ because $z_x^< + z_x^> = z_y^< + z_y^> = 4$ on the torus. The result therefore holds for any permutation on the torus.

Any permutation of the grid is obtained from some permutation on the torus by cutting between certain two rows and between certain two columns. Let π be a permutation on the torus, and let π' be a permutation on a cylinder obtained from π by cutting between two rows. Let x, y be two cells that are neighbors in π but not in π' . Let us denote by $b_1^<(x, y)$ the contribution of the cells x, y to $b_1^<$, and similarly for other quantities. We have $c_1'^<(x, y) = c_1^<(x, y) = 0 = c_1'^>(x, y) = c_1^>(x, y)$ because there are still no corners in π' . Suppose that $(\pi(x)_1, \pi(y)_1) \in S_1^< \times S_1^<$. Then $d_1'^>(x, y) = d_1^>(x, y)$, $b_1'^>(x, y) = b_1^>(x, y)$, while $d_1'^<(x, y) = d_1^<(x, y) - 1$ (because the small row defect between x and y has been

removed), and $b'_1{}^<(x, y) = b_1{}^<(x, y) + 2$ (because x, y are on the boundary of π'). Similarly if $(\pi(x)_1, \pi(y)_1) \in S_1^> \times S_1^>$. Suppose without loss of generality that $(\pi(x)_1, \pi(y)_1) \in S_1^< \times S_1^>$. Then $d'_1{}^<(x, y) = d_1{}^<(x, y)$, $d'_1{}^>(x, y) = d_1{}^>(x, y)$, $b'_1{}^<(x, y) = b_1{}^<(x, y) + 1$ and $b'_1{}^>(x, y) = b_1{}^>(x, y) + 1$. Similarly for all other pairs of cells across the cut, so the result holds for π' .

Finally, let π be a permutation on the cylinder, and let π' be obtained from π by cutting between two columns. Let x, y be two cells that are neighbors in π but not in π' . If x, y are not among the corners of π' then we can argue as above. Thus suppose that x, y are among the corners of π' , necessarily in the same row. Suppose that $(\pi(x)_1, \pi(y)_1) \in S_1^< \times S_1^<$. Then $c'_1{}^<(x, y) = c_1{}^<(x, y) + 2$, $b'_1{}^<(x, y) = b_1{}^<(x, y) - 2$ (because both x, y were on the boundary of π but are among the corners of π'), $d'_1{}^<(x, y) = d_1{}^<(x, y) - 1$, and the large values are not affected. Similarly if $(\pi(x)_1, \pi(y)_1) \in S_1^> \times S_1^>$. Suppose without loss of generality that $(\pi(x)_1, \pi(y)_1) \in S_1^< \times S_1^>$. Then $c'_1{}^<(x, y) = c_1{}^<(x, y) + 1$, $c'_1{}^>(x, y) = c_1{}^>(x, y) + 1$, $b'_1{}^<(x, y) = b_1{}^<(x, y) - 1$, $b'_1{}^>(x, y) = b_1{}^>(x, y) - 1$, and the defect values are not affected. Hence the result holds for π' , and we are through. \square

2.3. Separating the contributions of corners, boundaries and defects. Recall that maximizing $f(\pi)$ is equivalent to minimizing $g(\pi)$. The difficulty at hand is that the row and column coordinates cannot be optimized independently because the multisets B, C consist of values on cells, not of independent row and column values. Another difficulty is that the corners and defects are counted twice. In this subsection we derive a crucial inequality for $g(\pi)$ that removes most of the difficulties.

For a permutation π of $I(n_1, n_2)$, let

$$h(\pi) = \sum B_*^> + \sum C_*^> - \sum B_*^< - \sum C_*^<.$$

Furthermore, for $i \in \{1, 2\}$ let

$$x_i = b_i^< + c_i^< - b_i^> - c_i^>.$$

Proposition 2.6. *Let π be a permutation of $I(n_1, n_2)$ with $n_i = 2t_i$ and $n_1 \leq n_2$. Then*

$$g(\pi) \geq h(\pi) + 4 + n_1 + \sum_{i=1}^2 (t_i + 1/2)x_i.$$

Proof. By Proposition 2.5,

$$d_i^> - d_i^< = (b_i^< + 2c_i^< - b_i^> - 2c_i^>)/2 = x_i/2 + (c_i^< - c_i^>)/2,$$

and hence

$$\begin{aligned} \sum D_i^> - \sum D_i^< &\geq (t_i + 1)d_i^> - t_i d_i^< = (t_i + 1/2)(d_i^> - d_i^<) + (d_i^> + d_i^<)/2 \\ (2.2) \quad &= (t_i + 1/2)(d_i^> - d_i^<) + d_i/2 \\ &= (t_i + 1/2)x_i/2 + (t_i + 1/2)(c_i^< - c_i^>)/2 + d_i/2. \end{aligned}$$

Moreover,

$$\begin{aligned} (2.3) \quad &(t_i + 1/2)(c_i^< - c_i^>) + \sum C_i^> - \sum C_i^< \\ &= \left(\sum C_i^> - (t_i + 1/2)c_i^> \right) + \left((t_i + 1/2)c_i^< - \sum C_i^< \right) \\ &\geq c_i^>/2 + c_i^</2 = c_i/2 = 2. \end{aligned}$$

Using (2.2) and (2.3), we therefore have

$$\begin{aligned}
g(\pi) &= h(\pi) + \sum_{i=1}^2 \left(2 \sum D_i^> - 2 \sum D_i^< + \sum C_i^> - \sum C_i^< \right) \\
&\geq h(\pi) + \sum_{i=1}^2 \left((t_i + 1/2)x_i + (t_i + 1/2)(c_i^< - c_i^>) + \sum C_i^> - \sum C_i^< + d_i \right) \\
&\geq h(\pi) + \sum_{i=1}^2 \left((t_i + 1/2)x_i + 2 + d_i \right) \\
&= h(\pi) + 4 + d_* + \sum_{i=1}^2 (t_i + 1/2)x_i.
\end{aligned}$$

Finally, by Corollary 2.3 we have

$$(2.4) \quad d_* \geq n_1,$$

finishing the proof. □

2.4. Best approximations to disks. Let n_1, n_2 be positive integers, and let

$$O = ((n_1 + 1)/2, (n_2 + 1)/2)$$

be the geometric center of $I(n_1, n_2)$. For a subset A of $I(n_1, n_2)$ let

$$w(A) = \sum_{x \in A} d(x, O),$$

and for a positive integer k let

$$w_k(n_1, n_2) = \min\{w(A) : A \subseteq I(n_1, n_2), |A| = k\}.$$

A subset A of $I(n_1, n_2)$ is said to be a (*best approximation to a*) *disk* if $w(A) = w_{|A|}(n_1, n_2)$.

In order to understand disks in I , define the i th *layer* Λ_i by

$$\Lambda_i = \{x \in I(n_1, n_2) : d(x, O) = i\}.$$

Figure 1 depicts the layers $\Lambda_1, \Lambda_2, \Lambda_3$ in the situation when $n_1 = 4$ and $n_2 \geq 6$ is even.

	3	2	2	3	
3	2	1	1	2	3
3	2	1	1	2	3
	3	2	2	3	

FIGURE 1. The first three layers around the geometric center of $I(4, n_2)$.

Lemma 2.7. *A subset A of the grid $I(n_1, n_2)$ is a disk if and only if there are $r \geq 0$ and a subset P of Λ_{r+1} such that $A = P \cup \bigcup_{i=1}^r \Lambda_i$.*

Proof. Suppose that there are $i < j$, $x \in \Lambda_i$, $y \in \Lambda_j$ such that $x \notin A$ and $y \in A$. Then $A' = (A \cup \{x\}) \setminus \{y\}$ satisfies $|A'| = |A|$ and $w(A') = w(A) + i - j < w(A)$, a contradiction. The result follows. □

Before we return to the problem of neighbor separation, let us explicitly find the values $w_k(n_1, n_2)$ when $n_1 = 2t_1$, $n_2 = 2t_2$ are even and $n_1 \leq n_2$, using Lemma 2.7. The formula is straightforward but a bit cumbersome. First, we have

$$|\Lambda_i| = \begin{cases} 4i, & \text{if } 0 \leq i \leq t_1, \\ 2n_1, & \text{if } t_1 < i \leq t_2, \\ 2n_1 - 4(i - t_2) = 2(n_1 + n_2 - 2i), & \text{if } t_2 < i \leq t_1 + t_2, \\ 0, & \text{if } i > t_1 + t_2. \end{cases}$$

Given k such that $0 \leq k \leq n_1 n_2$, let r be the largest integer such that $\sum_{i=1}^r |\Lambda_i| \leq k$. Then, by Lemma 2.7,

$$(2.5) \quad w_k(n_1, n_2) = \sum_{i=1}^r i|\Lambda_i| + (r+1) \left(k - \sum_{i=1}^r |\Lambda_i| \right) = (r+1)k - \sum_{i=1}^r (r+1-i)|\Lambda_i|.$$

Here is the key connection between permutations and disks in $I(n_1, n_2)$:

Lemma 2.8. *Let π be a permutation of $I(n_1, n_2)$ with $n_i = 2t_i$. Then*

$$w(B \cup C) = h(\pi) + \sum_{i=1}^2 (t_i + 1/2)x_i.$$

Proof. Consider a pair of cells in $B \cup C$, one with a small row value, say $(t_1 - i, u)$, and one with a large row value, say $(t_1 + j, v)$. Then the contribution of the first coordinates of these two cells to $w(B \cup C)$ is $((t_1 + 1/2) - (t_1 - i)) + ((t_1 + j) - (t_1 + 1/2)) = i + j$. On the other hand, the analogous contribution to $h(\pi)$ is $t_1 + j - (t_1 - i) = i + j$. Similarly for small and large column values.

Now consider all cells of $B \cup C$. Pair together as many first coordinates of these cells as possible, subject to the constraint that small row values are matched with large row values. Pair column values similarly. For these matched pairs of coordinates, the contribution to $w(B \cup C)$ and $h(\pi)$ is the same, as we have just shown. If $x_1 \geq 0$, there are x_1 small row values remaining (unmatched), otherwise there are $-x_1$ large row values remaining. Similarly for columns.

Let the $|x_1| = q$ unmatched row values be i_1, \dots, i_q . If $x_1 \geq 0$, the contribution of these values to $h(\pi)$ is $-\sum_{j=1}^q i_j$, while the contribution to $w(B \cup C)$ is $\sum_{j=1}^q ((t_1 + 1/2) - i_j) = -\sum_{j=1}^q i_j + (t_1 + 1/2)q = -\sum_{j=1}^q i_j + (t_1 + 1/2)x_1$. If $x_1 \leq 0$, the contribution of these values to $h(\pi)$ is $\sum_{j=1}^q i_j$, while the contribution to $w(B \cup C)$ is $\sum_{j=1}^q (i_j - (t_1 + 1/2)) = \sum_{j=1}^q i_j + (t_1 + 1/2)x_1$. Similarly for columns, and the result follows. \square

Since $|B \cup C| = 2n_1 + 2n_2 - 4$, we deduce from Proposition 2.6 and Lemma 2.8:

Corollary 2.9. *Let π be a permutation of $I(n_1, n_2)$ with $n_i = 2t_i$ and $n_1 \leq n_2$. Then*

$$g(\pi) \geq 4 + n_1 + w(B \cup C).$$

In particular,

$$(2.6) \quad g(\pi) \geq 4 + n_1 + w_{2n_1+2n_2-4}(n_1, n_2).$$

3. OPTIMAL SOLUTIONS

In this section we construct a family of optimal solutions and prove the main result, Theorem 3.7.

3.1. A family of optimal solutions. Suppose that $2t_1 = n_1 \leq n_2 = 2t_2$. Note that the inequality (2.6) for π is in fact an equality if and only if all of the following conditions hold:

- (2.3) is an equality, that is, every value of $C_i^>$ is equal to $t_i + 1$ and every value of $C_i^<$ is equal to t_i , which is equivalent to $C = \Lambda_1$.
- (2.4) is an equality, that is, $d_* = n_1$. By Lemma 2.2, this is equivalent to the grid of π being evenly split into a homogeneous half and a heterogeneous half, with all defects occurring across the dividing line. In particular, $x_1 = x_2 = 0$. The actual placement of entries of Λ_1 into the corners determines the direction of the cut (we only have a choice if $n_1 = n_2$) and whether all defects will be row defects or column defects. (See Example 3.2.)
- (2.2) is an equality, that is, every value of $D_i^>$ is equal to $t_i + 1$ and every value of $D_i^<$ is equal to t_i . We observe that the defects must then be either all row defects alternating between t_1 and $t_1 + 1$, or all column defects alternating between t_2 and $t_2 + 1$, as the following diagram illustrates:

$$\begin{array}{ccc}
 \begin{array}{c|c}
 (<, <) & (<, >) \\
 (>, >) & (>, <) \\
 \vdots & \vdots
 \end{array} & \text{or} & \begin{array}{c|c}
 (<, <) & (>, <) \\
 (>, >) & (<, >) \\
 \vdots & \vdots
 \end{array}
 \end{array}$$

- $w(B \cup C) = w_{2n_1+2n_2-4}(n_1, n_2)$, that is, $B \cup C$ is a disk in $I(n_1, n_2)$.

Let us describe the family $\mathcal{P}(n_1, n_2)$ of permutations on $I(n_1, n_2)$ that satisfy all these conditions. It will be clear that $\mathcal{P}(n_1, n_2)$ is nonempty when n_1, n_2 are sufficiently large. We will give exact bounds on n_1, n_2 in Subsection 3.2.

Construction 3.1. Let $2t_1 = n_1 \leq n_2 = 2t_2$. The following procedure constructs all permutations σ of $\mathcal{P}(n_1, n_2)$ with a vertical defect line. (When $n_1 = n_2$, the members of $\mathcal{P}(n_1, n_2)$ with a horizontal defect line are obtained by a transposed construction. When $n_1 < n_2$, no members of $\mathcal{P}(n_1, n_2)$ have a horizontal defect line.)

- (i) Populate the four corners of σ with cells from Λ_1 so that one half (left or right) of the grid is homogeneous and the other half is heterogeneous. This will determine if all defects will be row defects or column defects.
- (ii) Populate the boundary with entries from Λ_2, Λ_3 , and so on, starting with Λ_2 , always filling in four entries at a time, using one entry from each $S_1^> \times S_2^>, S_1^> \times S_2^<, S_1^< \times S_2^>, S_1^< \times S_2^<$. In case of row defects, make sure that the two defects (in top and bottom row across the defect line) have values t_1 and $t_1 + 1$. In case of column defects, the defect values must be t_2 and $t_2 + 1$.
- (iii) Populate half of the interior cells along the defect line, one in each row, so that the row defects alternate values $t_1, t_1 + 1$ (or the column defects alternate values $t_2, t_2 + 1$).
- (iv) Populate the remaining cells arbitrarily, subject only to the restrictions imposed by the double checkerboard.

Let us illustrate Construction 3.1 with three examples that will be helpful in Subsection 3.2. In all examples the partial grids are labeled according to the steps in Construction 3.1. To save space and to improve legibility, we write i, j instead of (i, j) as the values of π .

Example 3.2. Let $n_1 = n_2 = 6$. The four grids in Figure 2 show how to construct a permutation of $\mathcal{P}(n_1, n_2)$ with a vertical defect line and row defects in four steps.

In step (i) we have made the choice to have a vertical defect line (by placing the homogeneous values in the same column rather than in the same row) and to have row defects (by not transposing the locations of the entries $(3, 4)$ and $(4, 3)$, say).




FIGURE 2. A permutation of the 6×6 grid maximizing the separation of neighbors with respect to the L_1 metric.

The following diagram corresponds to our choices of corner cells (+), boundary cells (-), cells that contribute a defect (\square), and cells that are both on the boundary and contribute a defect (\boxplus). Note that in the row defect resulting from the neighboring cells $(3, 2)$, $(3, 5)$ we have arbitrarily assigned $(3, 2)$ as the cell contributing the defect.

	1	2	3	4	5	6
1			-	-		
2		-	-	-	-	
3	\square	\boxplus	+	+	-	\square
4	\square	\boxplus	+	+	-	\square
5		-	-	-	-	
6			-	-		

Example 3.3. Let $n_1 = 4$ and $n_2 = 12$. Using the same notation as in Example 3.2, here is a diagram for a permutation in $\mathcal{P}(4, 12)$

	1	2	3	4	5	6	7	8	9	10	11	12
1				-	-	-		-	-	-		
2		□	⊠	-	-	+	+	-	-	-		
3		□	⊠	-	-	+	+	-	-	-		
4				-	-	-	-	-	-	-		

Figure 3 gives a partial grid of a permutation $\mathcal{P}(4, 12)$ obtained by steps (i)–(iii) of Construction 3.1. Step (iv) can be routinely completed.

2,6	3,8	1,6	4,7	2,5	3,9	3,3	1,7	3,5	2,8	4,6	2,7
4,8					2,2						3,4
2,4						3,2					1,8
3,7	1,5	3,10	1,4	4,9	2,3	2,10	4,4	1,9	4,5	2,9	3,6

FIGURE 3. A partial, routinely completable permutation of the 4×12 grid maximizing the separation of neighbors with respect to the L_1 metric.

Example 3.4. Let $n_1 = 10$ and $n_2 = 12$. Using the same notation as in Example 3.2, here is a diagram for a permutation in $\mathcal{P}(10, 12)$

	1	2	3	4	5	6	7	8	9	10	11	12
1												
2						-	-					
3					-	-	-	-				
4				-	-	-	-	-	-			
5	□	□	⊠	-	-	+	+	-	-	-	□	□
6	□	□	⊠	-	-	+	+	-	-	-	□	□
7				-	-	-	-	-	-			
8				-	-	-	-	-	-			
9						-	-	-				
10												

Figure 4 gives a partial grid of a permutation $\mathcal{P}(10, 12)$ obtained by steps (i)–(iii) of Constructions 3.1. Step (iv) can be routinely completed.

3.2. Existence of $\mathcal{P}(n_1, n_2)$. Let $n_1 \leq n_2$ be positive even integers. Recall that $\mathcal{P}(n_1, n_2)$ consists of all permutations π of $I(n_1, n_2)$ for which the inequality (2.6) is in fact an equality. The following lemma characterizes those parameters n_1, n_2 for which $\mathcal{P}(n_1, n_2) \neq \emptyset$.

Lemma 3.5. *Let $2t_1 = n_1 \leq n_2 = 2t_2$, and let r be the smallest integer for which $\Lambda = \bigcup_{i=1}^r \Lambda_i$ satisfies $|\Lambda| \geq |B \cup C|$. Then $\mathcal{P}(n_1, n_2) \neq \emptyset$ if and only if one of the following conditions holds:*

- (i) $|\Lambda| = |B \cup C|$ and $r \leq t_2 - (t_1 - 1)/2$,

5,6	6,8	5,5	7,7	4,6	6,9	6,3	5,8	6,5	4,7	7,6	5,7
7,8					5,2						6,4
5,4					6,11						5,9
8,7					5,1						7,5
4,5					6,12						4,8
6,10						5,11					8,6
3,6						6,2					3,7
7,9						5,12					7,4
4,4						6,1					5,10
6,7	3,5	8,8	2,6	9,7	5,3	2,7	9,6	3,8	8,5	4,9	6,6

FIGURE 4. A partial, routinely completable permutation of the 10×12 grid maximizing the separation of neighbors with respect to the L_1 metric.

(ii) $|\Lambda| > |B \cup C|$ and $r \leq t_2 - (t_1 - 3)/2$.

Proof. We will again only discuss permutations with a vertical defect line. In order to have $\mathcal{P}(n_1, n_2) \neq \emptyset$, $B \cup C$ must form a disk in $I(n_1, n_2)$. Note that our choice of r is the smallest possible so that $B \cup C$ can be chosen as a subset of layers $\Lambda_1, \dots, \Lambda_r$. Defects must be chosen either in rows $t_1, t_1 + 1$, or in columns $t_2, t_2 + 1$. Row defects can be selected whenever column defects can be selected (since $n_1 \leq n_2$). We will therefore focus on row defects. Then $\mathcal{P}(n_1, n_2) \neq \emptyset$ if and only if it is possible to select t_1 defects in row t_1 (one of which is in $B \cup C$), and t_1 defects in row $t_1 + 1$ (one of which is again in $B \cup C$).

Suppose that $n_1 = 2$. Then no defects outside of $B \cup C$ are needed. On the other hand, $\Lambda = B \cup C$ is the entire grid, $r = t_2$, $t_1 = 1$, so the inequality $r \leq t_2 - (t_1 - 1)/2$ is satisfied. We can therefore assume that $n_1 > 2$.

If $|\Lambda| = |B \cup C|$ (as in Examples 3.3 and 3.4), we must pick $n_1 - 2$ defects outside of Λ , so we demand $4(t_2 - r) \geq n_1 - 2$, which is equivalent to $r \leq t_2 - (t_1 - 1)/2$.

Suppose that $|\Lambda| > |B \cup C|$ (as in Example 3.2). Then there are additional four candidates for defects in Λ . Suppose that $n_1 = 4$. We can then always select the four needed defects in Λ and, on the other hand, $r \leq t_2 - (t_1 - 3)/2$ holds because $r \leq t_2$ and $t_1 = 2$. We can therefore assume that $n_1 > 4$, pick six defects in Λ , and we need to have $4(t_2 - r) \geq n_1 - 6$, which is equivalent to $r \leq t_2 - (t_1 - 3)/2$. \square

Proposition 3.6. *Let n_1, n_2 be positive even integers. Then $\mathcal{P}(n_1, n_2) \neq \emptyset$ except in the cases $n_1 = n_2 \in \{4, 8, 12, 16\}$.*

Proof. Let $n_i = 2t_i$, $t_2 = t_1 + u$, and let r be the smallest integer such that $\Lambda = \bigcup_{i=1}^r \Lambda_i$ satisfies $|\Lambda| \geq |B \cup C| = 4(t_1 + t_2 - 1) = 4(2t_1 + u - 1)$. Recall that $|\Lambda_i| = 4i$ if $1 \leq i \leq t_1$ and $|\Lambda_i| = 4t_1$ if $t_1 < i \leq t_2$.

Suppose that $t_1 = 1$. Then $r = t_2$ and the conditions of Lemma 3.5 are satisfied. We can therefore assume that $t_1 > 1$.

It turns out that $r \leq t_2$. It suffices to show that $\sum_{i=1}^{t_2} |\Lambda_i| \geq |B \cup C|$, which is equivalent to $\sum_{i=1}^{t_1} |\Lambda_i| + \sum_{i=t_1+1}^{t_2} |\Lambda_i| = 2t_1(t_1 + 1) + 4t_1(t_2 - t_1) \geq 4(t_1 + t_2 - 1)$, i.e., $2t_2(t_1 - 1) \geq (t_1 + 2)(t_1 - 1)$, $2t_2 \geq t_1 + 2$, which holds because $t_2 \geq t_1 > 1$.

Now suppose that $r > t_1$ and $t_2 < 3t_1/2 - 1/2$. Then $2t_1(t_1 + 1) \leq \sum_{i=1}^{r-1} |\Lambda_i| < |B \cup C| = 4(t_1 + t_2 - 1) < 4(t_1 + 3t_1/2 - 1/2 - 1) = 10t_1 - 6$, so $t_1^2 - 4t_1 + 3 < 0$ and $t_1 = 2$. Keeping $t_2 < 3t_1/2 + 1/2$ in mind, we only need to check Lemma 3.5 for the cases $(t_1, t_2) \in \{(2, 2), (2, 3)\}$, both of which pass.

Next suppose that $r > t_1$ and $t_2 \geq 3t_1/2 - 1/2$, the latter yielding $m = t_2 - (t_1 - 1)/2 \geq t_1$. If $r \leq m$ then the conditions of Lemma 3.5 are satisfied, so suppose that $r > m$. Then $2t_1(t_1 + 1) + 4t_1(m - t_1) = \sum_{i=1}^m |\Lambda_i| \leq \sum_{i=1}^{r-1} |\Lambda_i| < 4(t_1 + t_2 - 1)$, which is equivalent to $t_2(t_1 - 1) < (t_1 - 1)(t_1 + 1)$ and thus to $t_2 < t_1 + 1$. Hence $t_2 = t_1$, a contradiction with $r \leq t_2$.

Finally suppose that $r \leq t_1$. We have $|\Lambda| = 4(1 + 2 + \dots + r) = 2r(r + 1)$, and the positive root of the quadratic equation $2r(r + 1) = 4(2t_1 + u - 1)$ is $\rho = (\sqrt{16t_1 + 8u - 7} - 1)/2$. If $|\Lambda| = |B \cup C|$, we have $\rho = r$, and we need to satisfy the inequality $\rho \leq t_2 - (t_1 - 1)/2 = u + t_1/2 + 1/2$ of part (i) of Lemma 3.5. If $|\Lambda| < |B \cup C|$, we have $\rho < r \leq \rho + 1$, and the inequality $\rho + 1 \leq t_2 - (t_1 - 3)/2$ (which we note is the same as in the case $|\Lambda| = |B \cup C|$) is at least as strong as the inequality we need to satisfy in part (ii) of Lemma 3.5. Now, the inequality $\rho \leq u + t_1/2 + 1/2$ is equivalent to

$$t_1^2 + (4u - 12)t_1 + (4u^2 + 11) \geq 0,$$

and it fails only in the cases $(u = 1 \text{ and } t_1 = 4)$ and $(u = 0 \text{ and } t_1 \in [2, 10])$. Out of these, only the cases $t_1 = t_2 \in \{2, 4, 6, 8\}$ actually fail the conditions of Lemma 3.5. \square

3.3. Main result. We can now state and prove the main result. See the comments following Theorem 3.7 for the exceptional cases $n_1 = n_2 \in \{4, 8, 12, 16\}$.

Theorem 3.7. *Let $n_1 \leq n_2$ be positive even integers such that either $n_1 < n_2$ or $n_1 = n_2 \notin \{4, 8, 12, 16\}$. Let $I(n_1, n_2) = \{1, \dots, n_1\} \times \{1, \dots, n_2\}$ be the $n_1 \times n_2$ grid equipped with the L_1 metric $d((x_1, x_2), (y_1, y_2)) = |x_1 - y_1| + |x_2 - y_2|$. Let*

$$N = \{\{x, y\} \in I(n_1, n_2) \times I(n_1, n_2) : d(x, y) = 1\}$$

be the set of unit neighbors in $I(n_1, n_2)$. For a permutation π of $I(n_1, n_2)$, let

$$f(\pi) = \sum_{\{x, y\} \in N} d(\pi(x), \pi(y)).$$

Then the set $\mathcal{P}(n_1, n_2)$ of Construction 3.1 is not empty, the function f attains maximum at π if and only if $\pi \in \mathcal{P}(n_1, n_2)$, and this maximum is equal to

$$(3.1) \quad n_1 n_2 (n_1 + n_2) - 4 - n_1 - w_{2n_1 + 2n_2 - 4}(n_1, n_2),$$

where $w_k(n_1, n_2)$ is as in (2.5).

Proof. Let $n_i = 2t_i$. Recall that $f(\pi)$ is maximized if and only if $g(\pi)$ is minimized, and $f(\pi) = \sum S_*^> - \sum S_*^< - g(\pi)$ by Proposition 2.4. By Corollary 2.9, $g(\pi) \geq 4 + n_1 + w(B \cup C)$. The discussion preceding Construction 3.1 shows that equality is achieved in Corollary 2.9 if and only if $\pi \in \mathcal{P}(n_1, n_2)$, in which case $w(B \cup C) = w_{2n_1 + 2n_2 - 4}(n_1, n_2)$ because $B \cup C$ must be a disk in $I(n_1, n_2)$. By Proposition 3.6, $\mathcal{P}(n_1, n_2) \neq \emptyset$.

It remains to show that $\sum S_*^> - \sum S_*^< = n_1 n_2 (n_1 + n_2)$. We have

$$\sum S_1^> - \sum S_1^< = 4n_2((t_1 + 1) + (t_1 + 2) + \dots + (t_1 + t_1) - 1 - 2 - \dots - t_1) = 4n_2 t_1^2.$$

Similarly, $\sum S_2^> - \sum S_2^< = 4n_1 t_2^2$. Hence $\sum S_*^> - \sum S_*^< = 4n_2 t_1^2 + 4n_1 t_2^2 = n_1 n_2 (n_1 + n_2)$. \square

It follows from our results that in the four exceptional cases $n_1 = n_2 \in \{4, 8, 12, 16\}$, the maximum of f must be less than (3.1). We claim that the maximum is actually equal to

$$(3.2) \quad n_1 n_2 (n_1 + n_2) - 6 - n_1 - w_{2n_1+2n_2-4}(n_1, n_2),$$

that is, two less than in the generic case.

It is easy to construct permutations π of $I(n_1, n_2)$ for which (3.2) is attained, by modifying Construction 3.1. It is more difficult to prove that the maximum must drop by two compared to the generic case, not just by one. The details will be presented elsewhere [1].

4. COMMENTS AND OPEN PROBLEMS

4.1. Maximum separation of neighbors for permutations on the torus. By mimicking the proof of Lemma 2.2, we obtain:

Lemma 4.1. *Let $n_1 \leq n_2$ be even integers. Consider an $n_1 \times n_2$ grid on a torus with $n_1 n_2 / 2$ red and $n_1 n_2 / 2$ blue vertices. Then there are at least $2n_1$ red-blue edges.*

Suppose that there are precisely $2n_1$ red-blue edges. If $n_1 < n_2$ then the grid can be split into monochromatic connected components by two vertical cuts. If $n_1 = n_2$ then the grid can be split into monochromatic connected components by two vertical cuts or by two horizontal cuts.

Theorem 4.2. *Let $n_1 \leq n_2$ be positive even integers, $n_i = 2t_i$. Let X be the $n_1 \times n_2$ toroidal grid, and let d be the L_1 metric on X . A permutation π of X maximizes f of (1.1) if and only if it consists of an $S_1^< \times S_2^<$, $S_1^> \times S_2^>$ checkerboard adjacent to an $S_1^< \times S_2^>$, $S_1^> \times S_2^<$ checkerboard split by two vertical lines (or two horizontal lines if $n_1 = n_2$) so that either along all defect lines the row defects alternate between $t_1, t_1 + 1$, or along all defect lines the column defects alternate between $t_2, t_2 + 1$. Such permutations exist.*

Proof. Since there are no corners and boundaries on X , $f(\pi)$ is maximized if and only if $g(\pi) = 2(\sum D_*^> - \sum D_*^<)$ is minimized. Color homogeneous entries blue and heterogeneous entries red. Every red-blue edge corresponds to a defect. By Lemma 4.1, the number of red-blue edges is at least $2n_1$, and the bound $2n_1$ is attained only by two vertical dividing lines (or by two horizontal dividing lines when $n_1 = n_2$, a possibility we will discard from now on).

Additional defects are avoided if and only if the red cells form an $S_1^< \times S_2^<$, $S_1^> \times S_2^>$ checkerboard and the blue cells form an $S_1^< \times S_2^>$, $S_1^> \times S_2^<$ checkerboard. This can certainly be arranged on X .

Consider now the cells along one of the vertical defect lines. Suppose there is a row defect along this line. Then, as in the non-toroidal case, all defects along this line will be row defects. Moreover, we claim that all defects along the other defect line will also be row defects. Suppose we encounter the defect $(<, <)|(<, >)$ across the first defect line, in a given row. If t_2 is odd, we will then see $(<, <)|(<, >)$ across the second defect line, while if t_2 is even, we will see $(>, >)|(>, <)$ there, in the same row.

The rest is easy. □

Example 4.3. A permutation π on the toroidal 6×6 grid that maximizes $f(\pi)$, with two vertical defect lines:

(6, 4)	(3, 1)	(2, 4)	(6, 1)	(3, 4)	(2, 1)
(1, 1)	(5, 4)	(4, 1)	(1, 4)	(5, 1)	(4, 2)
(6, 5)	(3, 2)	(2, 5)	(6, 2)	(3, 5)	(2, 2)
(1, 2)	(5, 5)	(4, 2)	(1, 5)	(5, 2)	(4, 5)
(6, 6)	(3, 3)	(2, 6)	(6, 3)	(3, 6)	(2, 3)
(1, 3)	(5, 6)	(4, 3)	(1, 6)	(5, 3)	(4, 6)

4.2. Open problems. In [4], the neighbor-separation problem was solved for permutations on 1-dimensional intervals with respect to the L_1 metric:

Theorem 4.4 ([4, Theorem 3.1]). *Let $X = \{1, \dots, n\}$ and let $d(i, j) = |i - j|$. Then the maximum value of f defined by (1.1) is*

$$\begin{aligned} & (2t^2 - 1)/(2t - 1), & \text{if } n = 2t, \text{ and} \\ & (2t^2 + 2t - 1)/(2t), & \text{if } n = 2t + 1. \end{aligned}$$

When $n = 2t$, the maximum is attained by precisely those permutations π that oscillate between $\{1, \dots, t\}$, $\{t + 1, \dots, n\}$ and satisfy $(\pi(1), \pi(n)) \in \{(t, t + 1), (t + 1, t)\}$.

When $n = 2t + 1$, the maximum is attained precisely by those permutations π that oscillate between $\{1, \dots, t\}$, $\{t + 1, \dots, n\}$ and satisfy $(\pi(1), \pi(n)) \in \{(t + 1, t + 2), (t + 2, t + 1)\}$, and by those that oscillate between $\{1, \dots, t + 1\}$, $\{t + 2, \dots, n\}$ and satisfy $(\pi(1), \pi(n)) \in \{(t, t + 1), (t + 1, t)\}$.

We found the 2-dimensional case in which at least one of the lengths is odd difficult to handle and we therefore ask:

Problem 4.5. *Let n_1, n_2 be positive integers, not both even. Describe the permutations π on $I(n_1, n_2)$ for which $f(\pi)$ is maximized with respect to the L_1 metric.*

The situation will be more intricate for 3-dimensional intervals. When n_1, n_2, n_3 are even and approximately the same, $f(\pi)$ appears to be maximized by a triple checkerboard with defects occurring along two perpendicular planes through the center of the prism. However, when n_1 is much bigger than n_2 and n_3 , $f(\pi)$ appears to be maximized by a triple checkerboard with defects occurring along two or more planes perpendicular to the first dimension.

Problem 4.6. *Let n_1, n_2, \dots, n_k be positive integers. Describe the permutations π of the $n_1 \times n_2 \times \dots \times n_k$ grid that maximize $f(\pi)$ with respect to the L_1 metric.*

ACKNOWLEDGMENT

We thank Dan Daly for pointing out several relevant references.

REFERENCES

- [1] Mohammed Albow, PhD Dissertation, in preparation.
- [2] Andreas Bärtschi, Barbara Geissmann, Daniel Graf, Tomas Hruz, Paolo Penna and Thomas Tschager, *On computing the total displacement number via weighted Motzkin paths*, In: Mäkinen V., Puglisi S., Salmela L. (eds) Combinatorial Algorithms. IWOCA 2016. Lecture Notes in Computer Science, vol 9843. Springer, Cham.

- [3] C. Berrou, Y. Saouter, C. Douillard, S. Kerouedan and M. Jezequel, *Designing good permutations for turbo codes: towards a single model*, 2004 IEEE International Conference on Communications (IEEE Cat. No.04CH37577), Paris, France, 2004, pp. 341–345.
- [4] Dan Daly and Petr Vojtěchovský, *How permutations displace points and stretch intervals*, *Ars Combinatoria* **90** (2009), 175–191.
- [5] Persi Diaconis and R.L. Graham, *Spearman’s footrule as a measure of disarray*, *J. Roy. Statist. Soc. Ser. B* **39** (1977), no. **2**, 262–268.
- [6] Roberto Gallero, Guido Montorsi, Sergio Benedetto and Giovanni Cancellieri, *Interleaver properties and their applications to the trellis complexity analysis of turbo codes*, *IEEE Transactions on communications* **49** (May 2001), no. **5**, 793–807.
- [7] Mathieu Guay-Paquet and T. Kyle Petersen, *The generating function for total displacement*, *Electron. J. Combin.* **21** (2014), no. **3**, Paper 3.37, 13 pp.
- [8] Panawit Hanpinitasak, Chalie Charoenlarnnoppaart and Prapun Suksompong, *2D interleaver design for image transmission over severe burst-error environment*, *The ECTI Transactions on Electrical Engineering, Electronics and Communications* **12**, no *2* (2014), 55–63.
- [9] Donald E. Knuth, *The Art of Computare Progamming, Volume 3, Sorting and Searching*, Second Edition, Addison-Wesley 1998.

(Albow, Edgington, Lopez) DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF DENVER, 2155 EAST WESLEY AVENUE, DENVER, COLORADO 80208, U.S.A.

E-mail address, Albow: Mohammed.Al-Bow@du.edu

E-mail address, Edgington: Jeffrey.Edgington@du.edu

E-mail address, Lopez: mlopez@du.edu

(Vojtěchovský) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DENVER, 2390 S YORK STREET, DENVER, COLORADO 80208, U.S.A.

E-mail address, Vojtěchovský: petr@math.du.edu