

Neighborhood Unions and Independent Cycles

Ronald J. Gould* Kazuhide Hirohata† Paul Horn‡

Abstract

In this paper, we investigate sufficient conditions on the neighborhood of independent vertices which imply that a graph contains k independent cycles or k independent chorded cycles. This is related to several results of Corrádi and Hajnal, Justesen, Wang, and Faudree and Gould on graphs containing k independent cycles, and Finkel on graphs containing k chorded cycles. In particular, we establish that if every pair of independent vertices in G have neighborhood union at least $2k + 1$ then G has k independent cycles, as long as $|G| > 30k$. This settles a conjecture of, and improves a result due to Faudree and Gould, who establish that $3k$ suffices. Additionally, we show that a graph having neighborhood union of every pair of independent vertices at least $4k + 1$ has at least k independent chorded cycles; Finkel previously established that minimum degree $3k$ is also a sufficient condition for this.

In 1963, Corrádi and Hajnal [1] verified a conjecture of Erdős by showing that every graph G on at least $3k$ vertices with minimum degree $2k$ contains k independent cycles. In the nearly fifty years since then, this result has been generalized in a number of different ways.

In 1989, Justesen [4] showed that the minimum degree condition can be replaced by a condition on the sum of degrees of non-adjacent vertices. This was improved to a sharp result by Wang [5] who showed the condition that the minimum degree was at least $2k$ could be replaced by the condition that

*Department of Mathematics and Computer Science, Emory University, Atlanta, GA.
Email: rg@mathcs.emory.edu

†Department of Electronic and Computer Engineering, Ibaraki National College of Technology, Ibaraki, Japan. Email: hirohata@ece.ibaraki-ct.ac.jp

‡Department of Mathematics and Computer Science, Emory University, Atlanta, GA.
Email: phorn@mathcs.emory.edu

the sum of degrees of non-adjacent vertices was at least $4k - 1$. Corrádi and Hajnal's result was generalized in a different direction by Finkel [3] who showed a graph on at least $4k$ vertices with minimum degree at least $3k$ contains at least k independent chorded cycles.

In a different direction, J. Faudree and Gould [2] showed that the condition on minimum degrees in the Corrádi-Hajnal theorem could be replaced by a condition on the neighborhood of non-adjacent vertices. In particular they showed that if G is a graph on at least $3k$ vertices such that the neighborhood of any two non-adjacent vertices has size at least $3k$, then G contains k independent cycles. This result is sharp in the sense that requiring a neighborhood of size $3k - 1$ is insufficient for $k = 1$, but it was conjectured it might be possible to replace their condition by one of the form $2k + O(1)$.

In this paper, we consider both the problem of finding independent cycles and independent chorded cycles in graphs under conditions on the neighborhood union of pairs of non-adjacent vertices. Before we state the main theorems, let us briefly introduce some notation. We indicate by $N(x)$ the set of neighbors of x , and $N[x] = N(x) \cup \{x\}$ denotes the closed neighborhood. For a set $S \subseteq V$, we denote by $N_S(x) = N(x) \cap S$. For a pair of non-adjacent vertices x, y we let $N(x, y) = N(x) \cup N(y)$, and define $N_S(x, y)$ in the obvious fashion. We define $x \sim y$ to mean that x is adjacent to y .

We begin by generalizing the result of Finkel as follows:

Theorem 1. *Suppose G is a graph on at least $4k$ vertices, such that any pair of non-adjacent vertices x, y has $|N(x, y)| \geq 4k + 1$. Then there exist k independent chorded cycles in G .*

Theorem 1 is sharp in the sense that it does not hold if the condition $|N(x, y)| \geq 4k + 1$ is replaced by $|N(x, y)| \geq 4k$. Here it fails even when $k = 1$: consider a collection of t independent K_3 's; then the neighborhood of any two non-adjacent vertices is 4, but there are no chorded cycles. If one prefers a connected counterexample, one may append a different K_3 to each vertex of a path or cycle.

In addition, we also prove the conjecture of Faudree and Gould [2]:

Theorem 2. *If G is a graph on more than $30k$ vertices, such that any pair of non-adjacent vertices x, y has $|N(x, y)| \geq 2k + 1$, then G contains k independent cycles.*

This result is also sharp, in the sense that $2k + 1$ cannot be relaxed to $2k$. Examples when $k = 1$ include disjoint edges, or trees where all leaves are adjacent to vertices of degree 2. For $k = 2$, the wheel graph serves as an example: non-adjacent vertices have neighborhood union at least four, but there are no two independent cycles.

It is possible that the constant '30' in the theorem can be improved, possibly even to 4. Indeed, one may slightly improve this constant by being more careful in a few places in our proof, but it is clear that such improvements will not bring the constant all the way to 4. Such an improvement seems to require new ideas.

1 The Proof of Theorem 1

Before we delve into the proof, we need a definition and some examples. Since graphs with minimum degree at least 3 contain a chorded cycle, we will often be concerned with graphs that contain no subgraph with minimum degree at least 3. A graph H of order h is called *2-degenerate* if it has the property that there is an ordering of vertices v_1, v_2, \dots, v_h of H such that each vertex has at most two edges to higher labeled vertices. We call such an ordering a *2-degenerate ordering*. Any graph containing no subgraph with minimum degree at least 3 is easily seen to be 2-degenerate.

We say that H contains a *k-tower* if there is a 2-degenerate ordering of G , say v_1, \dots, v_h , such that if v_i is the first vertex at distance $k - 1$ from v_1 , then for $2 \leq j \leq i$, each intermediate vertex v_j is adjacent to a vertex v_ℓ with $\ell < j$, that is, each v_j has an edge to the left in the ordering. One interpretation of this definition is that there is a 2-degenerate ordering starting at some vertex v such that a vertex at distance $k - 1$ from v can be lowered to degree 2 just by iteratively deleting vertices of degree 2 exposed by deleting v and its neighbors (and iterative neighbors) while leaving other vertices of degree 2 in the graph.

One may consider this definition an analogue of having a path of order k in a tree starting at a leaf and not containing any vertices of degree larger than 2. Clearly any 2-degenerate graph H must contain a 1-tower (this is simply a vertex of degree at most 2), and if H contains a k -tower, it clearly also contains a $(k - 1)$ -tower. A star is an example of a graph which only contains a 1-tower, while a path H would be an $|H|$ -tower. Also,

a 2-degenerate graph H which has a single vertex of degree 2 contains a $(\text{diam}(H) + 1)$ -tower.

Proof of Theorem 1. The proof is by induction on k and let $n = |G|$. For $k = 1$, and any n the result is clear: either $\delta(G) \geq 3$ and the result holds by the theorem of Finkel, or there exists a vertex x with $|N(x)| \leq 2$. If x is a vertex of degree ≤ 2 , since $|N(x, y)| \geq 5$ for all $y \not\sim x$, note that $N_{G-N[x]}(y) \geq 3$. Since $\delta(G - N[x]) \geq 3$, there must be a chorded cycle in $G - N[x]$.

We now suppose that the theorem holds for all $k' < k$, and show it holds for k . Let G be a graph satisfying the hypothesis of Theorem 1. If there exists a chorded 4-cycle C in G , then note that $G - C$ satisfies the hypothesis of Theorem 1 for $k' = k - 1$. Thus, by the inductive hypothesis, there exists $k - 1$ independent chorded cycles in $G - C$ and, along with C , we have k independent chorded cycles. Thus we may assume that G contains no chorded 4-cycles.

Note that G satisfies the conditions for $k' = k - 1$ (even after removing 4 arbitrary vertices), and hence by the inductive hypothesis there exist $k - 1$ independent chorded cycles in G : C_1, C_2, \dots, C_{k-1} . Let $\mathcal{C} = \bigcup C_i$. We shall assume that $|\mathcal{C}|$ is minimized. Let $H = G \setminus \mathcal{C}$, and assume that H has the maximum number of edges (subject to the constraint that $|\mathcal{C}|$ is minimized). If H has a chorded cycle we are done, so we may assume that H does not contain a chorded cycle and derive a contradiction. We note that $|H| \geq 4$. By the results of Finkel, the fact that H has no chorded cycles implies that H is 2-degenerate. We shall assume (subject to having the maximum number of edges) that H has a k -tower for as large a k as possible.

We make a few initial observations about H :

Claim 1. *If there is a $C_i \in \mathcal{C}$ and $x \in H$ with $|N_{C_i}(x)| \geq 3$, then $|C_i| \leq 6$.*

Proof. Suppose x is adjacent to $y_1, y_2, y_3 \in C_i$. If the distance between y_i and y_j on C is greater than 2 for any i, j , then there is a shorter chorded cycle using x (possibly using the edge to the other y_k as the chord.), contradicting the minimality of $|\mathcal{C}|$. Furthermore, if y_1, y_2, y_3 are a strip of 3 adjacent vertices in C_i , then they form a chorded 4-cycle with x , and we are done. The only way, then, that all of the pairwise distances of y_1, y_2, y_3 can be at most 2 is if $|C_i| \leq 6$. \square

Claim 2. *If $x \not\sim y$ with $|N_H(x, y)| \leq 4$ and $x, y \in H$, then there exists a chorded 6-cycle $C_i \in \mathcal{C}$ such that, up to relabeling x and y are adjacent to C_i in one of the following ways (see Figure 1):*




Figure 1

Proof. Note that $|N_{\mathcal{C}}(x, y)| \geq 4k+1-4 = 4k-3$, so there exists some $C_i \in \mathcal{C}$ such that $|N_{C_i}(x, y)| \geq 5$. Without loss of generality, say, $|N_{C_i}(x)| \geq 3$, and hence by Claim 1, $|C_i| \leq 6$.

If $|C_i| = 5$, and $|N_{C_i}(x)| \geq 4$, then there is a chorded 4-cycle in $C_i \cup \{x\}$. Thus $|N_{C_i}(x)| = 3$, and $N_{C_i}(x, y) = C_i$. It is easy to check that however x and y send edges into C_i , they would create a chorded 4-cycle, therefore $|C_i| = 6$.

If the chord in C_i connects two vertices at distance 2 on the cycle, then however the 3 edges from x are placed in C_i so that the ends have pairwise distance 2 on the cycle, they admit a shorter chorded cycle. (Recall, these neighbors must be at distance 2 on the cycle.) If both ends of the chord are adjacent to x , then there is a chorded 4-cycle. Otherwise there is a chorded 5-cycle. Therefore the chord bisects the C_6 as in the Figure 1. Then $|N_{C_i}(x)| = 3$, or else there would be a chorded 5 cycle; and since the edges must be pairwise at distance 2 along the cycle they must lie as in Figure 1. Then y has at least two neighbors in C_i which are not neighbors of x , as in Figure 1. \square

A corollary of Claim 2 is the following swapping lemma, which is key to our argument.

Lemma 1 (Swapping Lemma). *Suppose x, y are non-adjacent vertices in H with $|N_H(x, y)| \leq 4$. Then there exists a $C_i \in \mathcal{C}$ and vertices $z_x, z_y \in C_i$ such that $z_x \sim x$, $z_y \sim y$ and both $(C_i - \{z_x\}) \cup \{y\}$ and $(C_i - \{z_y\}) \cup \{x\}$*

are chorded cycles. Furthermore, for $x', y' \in V(H)$, if $x' \sim x$ then $z_x \not\sim x'$, likewise if $y' \sim y$ then $z_y \not\sim y'$.

Proof. By Claim 2, we know the adjacencies of x and y to some C_i . It is easy to verify, by inspection, that a chorded cycle including, say, x may be routed excluding one of the corner vertices adjacent to y ; that corner vertex will be denoted z_y . The lemma easily follows from inspection; a typical case is as in Figure 2:




Figure 2

Here the outer cycle of the new chorded cycle is highlighted. The assertion that $z_x \not\sim x'$ follows from the fact that, if it is adjacent, that it would create a chorded 5-cycle. This is easy to check, details are included as part of the proof of Lemma 2 below. \square

As an application of the swapping lemma, we see that (if we do not insist that the number of edges of H is maximized) we may assume H is connected: Suppose H is such that it has a minimum number of components, and that the largest component is as large as possible. If H had two components, say X_1 and X_2 where X_1 is the largest component, each would have a vertex of degree at most 2 in H . Suppose $x_1 \in X_1$ and $x_2 \in X_2$ are these vertices; then $|N_H(x_1, x_2)| \leq 4$. Applying the swapping lemma says that we may remove x_2 and append a vertex to x_1 , which would increase the size of the largest component, a contradiction. Thus H must be connected. In particular, when the number of edges in H is maximized, we know that $|E(H)| \geq |H| - 1$. Also note that if the number of edges in H is maximized, and we apply the swapping lemma we know that $|N_{H-\{y\}}(z_x)| \leq |N_H(y)|$ and similarly for z_y and x . The swapping lemma also implies that there are no isolated vertices in H .

Observe the following: Suppose k is the largest integer such that H has a k -tower. If $k \leq 2$, then G has at least 2 non-adjacent vertices of degree at

most 2. Indeed, if H has only a single vertex of degree ≤ 2 , then it has a $(\text{diam}(H) + 1)$ -tower, and since $|H| \geq 4$, $\text{diam}(H) \geq 2$. If H only has two vertices of degree ≤ 2 , and they are adjacent, the same conclusion holds. H cannot have just 3 pairwise adjacent vertices of degree ≤ 2 : if H had such, they would form one component of size 3, but $|H| \geq 4$, so there must be another component which must contain a vertex of degree ≤ 2 . Furthermore, if H has only a one-tower, and has only two non-adjacent vertices of degree ≤ 2 , then they must both be connected to a vertex of degree at most 4.

The following is the crux of the proof:

Lemma 2. *Suppose x, y, z are three vertices in H with $x \sim y$, $y \sim z$ and $x \not\sim z$, such that $|N_C(x, z)| + |N_C(y)| \geq 6k - 5$. Then $|\mathcal{C}|$ is not minimized.*

Proof. Suppose that $|\mathcal{C}|$ is minimized. We shall find a shorter chorded cycle, and thus a contradiction.

By hypothesis, there exists some chorded cycle $C = C_i \in \mathcal{C}$ such that $|N_C(x, z)| + |N_C(y)| \geq 7$. Clearly one of x, y , or z sends at least 3 edges into C , so by Claim 1, $|C| \leq 6$. If any one vertex sends 4 edges into C , it is clear that there is a shorter chorded cycle. We consider the other scenarios:

If $|N_C(y)| = 3$, then $|N_C(x, z)| \geq 4$. First, consider $|C| = 5$. Since there is only one way to fit 3 edges from a single vertex into a chorded 5-cycle without creating a shorter chorded cycle, we know the adjacencies from y to C . If one of x or z (say x) also has degree 3 in C , then it is easily seen that $N_C(x) = N_C(y)$. Then there are clearly chorded 4-cycles. Now suppose x and z both have degree 2 in C . Suppose b_1, b_2 are the two vertices which lie on one side of the chord. Both $b_1, b_2 \in N(y)$. One of b_1, b_2 (say b_1) is in the neighborhood of one of x or z (say x). But then we have a chorded 4-cycle: x, b_1, b_2, y with chord $\{y, b_1\}$.

Thus we may assume $|C| = 6$. If C is a chorded 6-cycle with a single vertex a on one side of the chord, and three vertices b_1, b_2, b_3 on the other (where b_1, b_3 are at distance 2 from a , and b_2 is at distance 3 and b_0 and b_4 are the remaining vertices according to this ordering), then if y has two consecutive neighbors on C or is adjacent to both ends of the chord, a shorter chorded cycle exists. Then the only possibility is that $y \sim \{a, b_1, b_3\}$. But then y, b_3, b_4, b_0, a is a chorded 5-cycle with chord ab_4 . Thus, C must be the chorded 6-cycle as in Figure 2. Suppose the vertices of the chord, arranged in order along the cycle are x_1, \dots, x_6 , and the chord is $\{x_1, x_4\}$. Then $y \sim \{x_1, x_3, x_5\}$. If $x \sim x_1$ (or $z \sim x_1$) then there is a chorded 5-cycle:

x, x_1, x_2, x_3, y with chord yx_1 , so $x \not\sim x_1$ and similarly $x \not\sim \{x_3, x_5\}$. But $|N_C(x, z)| \geq 4$ so clearly x or z must be adjacent to one of those vertices. Thus $|N_C(y)| < 3$.

If $|N_C(y)| = 2$, then $|N_C(x, z)| \geq 5$. As argued in Claim 2, $|C| = 6$, and it is of the form described above: vertices x_1, \dots, x_6 with chord x_1x_4 where (without loss of generality) $x \sim \{x_1, x_3, x_5\}$. Furthermore, either $z \sim \{x_2, x_4\}$ or $z \sim \{x_2, x_6\}$. First consider the case where $z \sim \{x_2, x_4\}$ (as the other case is similar, but with chord from x to C). Then if $y \sim x_1$ we have a chorded 5-cycle x, x_1, x_2, z, y with chord yx_1 . Likewise y cannot be adjacent to any vertex in $N_C(x, z)$. But $|N_C(y)| = 2$, so this must be the case. (Here the key property is that chorded cycles are formed using exactly one edge in C .) Similarly if $|N_C(y)| = 1$, and $|N_C(x, z)| = 6$; the fact that $N_C(y) \in N_C(x, z)$ as $|C| = 6$ will force a similar chorded 5-cycle, using x, y, z and one edge in C , with the edge from y or x to C acting as the chord. \square

The remainder of the proof of Theorem 2 is to show that by the Swapping Lemma, Lemma 1, we may ensure that H has vertices x, y, z that satisfy the conditions of Lemma 2. This leads to a contradiction to the assumption that H had no chorded cycles, completing the proof. We now consider the cases:

First suppose that H has no 3-tower. Thus it has two non-adjacent vertices x and y of degree at most 2. Let us assume that x is at the start of the highest tower in H (which by assumption is at most a 2-tower). First let us assume that there is a 2-tower starting at x . We apply the swapping lemma to exchange y for a $y' \sim x$. Let C' and H' denote the new cycle system and new copy of H after the swap. Note that $|N_{H'}(y')| \leq 2$ and y' is adjacent to no neighbors of x as assertions of the swapping lemma. Thus, there is a 3-tower starting at y' and by the maximality of the tower height, this implies that $|N_{H'}(y')| \leq 1$. Now since there is a 3-tower starting at y' there is a neighbor $x' \sim x$ such that $x' \not\sim y'$ and $|N_{H'}(y', x')| \leq 3$ (this is the vertex which has degree at most 2 after deletion of y' and x). Now one of y' and x' (say x') has $|N_{C'}(x')| > 2k - 1$ by the neighborhood condition. Apply the swapping lemma to exchange y' for a $z \sim x'$ (if y' is the vertex of high degree, do this in reverse). Then if H'' is the resulting leftover vertices $|N_{H''}(z)| \leq 2$, and $z \not\sim x$ and it is easy to see that z, x', x satisfy the conditions of Lemma 2 as $|N_{H''}(z, x)| \leq 4$ so $|N_{C''}(z, x)| \geq 4k + 1 - 4 = 4k - 3$ and $|N_{C''}(x')| \geq 2k - 2$, where C'' is the new cycle set after the swaps.

If there is a 1-tower, but no 2-tower the proof is similar. Again, we use H' , H'' , etc. to denote the set H after swaps, and \mathcal{C}' accordingly. We may apply the swapping lemma to find $y' \sim x$, there will be a 2-tower starting at y' and hence $|N_{H'}(y')| = 1$. It is possible that, in fact, a 3-tower now starts at y' . In this case there is a vertex $x' \sim x$ such that $|N_{H'}(x', y')| \leq 3$ and we are in the same case as before. If not, then there is a vertex z of degree at most 2 non-adjacent to y' . We may again apply the swapping lemma to find a neighbor $z' \sim y'$ with $z' \not\sim x$ and $|N_{H''}(z')| \leq 2$. Now, $|N_{H''}(z', x)| \leq 4$ (as their neighborhoods contain the common vertex y') and we are again in the position above: one of z' or x (say x) has high degree to \mathcal{C}'' , we apply the swapping lemma to find a $z'' \sim x$ and end up in a situation where the hypothesis of Lemma 2 are satisfied.

Thus we may assume that H has a 3-tower. Suppose x is the degree (at most) 2 vertex first in the ordering defining the tower. Then there is a vertex y at distance 2 from x with $|N_H(x, y)| \leq 4$.

First suppose that there exists such a y with $|N_H(y)| \leq 3$, and let z denote the center vertex between x and y .

Suppose $|N_H(y)| \leq 3$. As $|N_{\mathcal{C}}(x, y)| \geq 4k - 3$, one of x or y has neighborhood in \mathcal{C} of size at least $2k - 1$. If this is x , first consider the case where there is another vertex of degree at most 2 in H , say w . If so, applying the swapping lemma to find $w' \sim x$ and taking a vertex z' of degree 3 adjacent to x gives a set w, x, z' which satisfies the conditions of Lemma 2. If this is not x , we consider separately the cases where $N_H(x)$ is connected and disconnected. If $N_H(x)$ is connected we see that we can choose $z \sim y$ with $N_H(z) = \{x, y, z'\}$, where z' and z are the neighbors of x . Then after applying the swapping lemma to y and x to find $y' \sim x$ with $|N_H(y')| \leq 3$, we may assume that $|N_{H-y}(z)| = 2$. In particular $|N_H(z, y')| \leq 4$ (as the neighborhoods share x) and z, x, y' satisfy the conditions of Lemma 2. If $N_H(x)$ is not connected, the fact that $|N_H(y)| \leq 3$ says we may choose y such that a middle vertex z has $|N_{H-y}(z)| \leq 2$. Again, apply the swapping lemma to find $y' \sim x$ with $|N_H(y')| \leq 3$ and y', x, z meeting the conditions of Lemma 2. If y has the high degree instead, we apply the Swapping Lemma to remove x and find $x' \sim y$, and x', y, z will similarly meet the criteria of Lemma 2.

If $|N_H(y)| \geq 4$ for all y at distance 2 from x , there is one final case. Here x is adjacent to two vertices of degree 3, x_1, x_2 , and x_1 and x_2 are both adjacent to the same two vertices y_1, y_2 (one of which must have degree at

most 4). Let y the vertex of degree 4. Note that $|N_H(x_1, x_2)| = 3$, so one of x_1, x_2 (say x_1) has $|N_C(x_1)| \geq 2k - 1$. Apply the switching lemma to x_1 and x_2 to find a vertex $x'_2 \sim x_1$ with $|N_H(x'_2)| \leq 3$ and $x'_2 \not\sim x$. Then $|N_{H-x_2}(x)| = 1$, and $|N_C(x_1)| \geq 2k - 2$. Thus $|N_H(x'_2, x)| \leq 3$, and x'_2, x_1, x satisfy the conditions of Lemma 2.

In all cases, we have reached a contradiction, hence H must have contained a chorded cycle and we have completed the proof of Theorem 1. \square

2 The Proof of Theorem 2

Like the proof of Theorem 1, the general strategy will be by induction on k , for any fixed $n = |G|$. The case $k = 1$ is trivial: if G has minimum degree at least 2, clearly G contains a cycle; otherwise there is a vertex x of degree 1, and $G - N[x]$ has minimum degree ≥ 2 by the neighborhood condition, and thus contains a cycle.

We say that a sequence v_1, \dots, v_k is a *non-branching path* of order k in G , if $v_i \sim v_{i+1}$, $1 \leq i \leq k - 1$, and $|N(v_i)| \leq 2$ for all $i \leq k$.

Suppose G is a graph satisfying the hypothesis of Theorem 2 for some k , and that the theorem has been proved for $k - 1$. The hypothesis of the theorem are satisfied for $k - 1$, so G contains $k - 1$ independent cycles C_1, \dots, C_{k-1} . Let $\mathcal{C} = \bigcup_{i=1}^{k-1} C_i$ and assume that $|\mathcal{C}|$ is minimized. Let $H = G - \mathcal{C}$. We assume H has a minimum number of components. If H contains a cycle, then we have exhibited k disjoint cycles and are done, so we may assume H is a forest. We prove the following swapping lemma, similar to Lemma 1: Here, H' is the graph induced by $(H - \{x_i\}) \cup z$.

Lemma 3. *Suppose x_1, x_2 , and x_3 are vertices in H which are pairwise non-adjacent and $|N_H(x_i)| \leq 1$. Then there exist $i, j \in \{1, 2, 3\}$ with $i \neq j$, $C_k \subseteq \mathcal{C}$ and $z \in C_k$, such that $(C_k - \{z\}) \cup \{x_i\}$ is a cycle and $z \sim x_j$. Furthermore $|N_{H'}(z)| = 1$.*

Remark: There is one key difference between this lemma and Lemma 1. In Lemma 1 we are able to choose whether to append the new vertex to x or append the new vertex to y . Here, we have no choice: the lemma guarantees that we can swap some x_i for some z adjacent to some x_j , but we cannot guarantee (say) that we can add x_1 to a cycle and swap it out for a $z \sim x_2$.

To prove Lemma 3 we make the following observation, which follows immediately from the minimality of $|\mathcal{C}|$:

Claim 3. *Suppose $x \in H$ has $|N_{C_i}(x)| \geq 2$, then $|C_i| \leq 4$. Furthermore, if $|N_{C_i}(x)| = 3$, then $|C_i| = 3$.*

Proof of Lemma 3. Note that

$$|N_{\mathcal{C}}(x_1, x_2)| + |N_{\mathcal{C}}(x_1, x_3)| + |N_{\mathcal{C}}(x_2, x_3)| \geq 3(2k - 1)$$

so there exists a cycle C_i such that

$$|N_{C_i}(x_1, x_2)| + |N_{C_i}(x_1, x_3)| + |N_{C_i}(x_2, x_3)| \geq 7. \quad (1)$$

In particular, for some pair x_a, x_b (say x_1 and x_2) we have $|N_{C_i}(x_1, x_2)| \geq 3$. Thus $|C_i| \leq 4$ by Claim 3. At least one of x_1 and x_2 have degree at least 2 into C_i , say x_1 . If $|N_{C_i}(x_1)| = 2$, then we will set $x_a = x_1$ and $x_b = x_2$ and one can easily check that the conditions of the lemma are satisfied, since x_2 must have a neighbor in C_i that is not amongst the neighbors of x_1 . If $|N_{C_i}(x_1)| = 3$, then $|C_i| = 3$. In this case, note that one of x_2 or x_3 must have a neighbor in C_i , otherwise (1) will not hold; if this is x_3 , say, let $x_a = x_1$, and $x_b = x_3$. The other cases follow analogously. \square

Note that Lemma 3 implies the following:

Claim 4. *H has at most 2 isolated vertices.*

Proof. Otherwise, applying the swapping lemma (Lemma 3) to these vertices would reduce the number of components. \square

The key observation is the following:

Lemma 4. *Without loss of generality, we may assume that H contains either two disjoint non-branching paths of order at least 4, one of which starts at a leaf or a single non-branching path of order 8, which need not contain a leaf.*

Proof. For the purposes of this, we assume that H has no isolated vertices; even after removing the isolated vertices, we decrease the order of H by at most 2 by Claim 4. We assume that H cannot have a non-branching path

of order 8 starting at a leaf, subject to the conditions imposed on H . Under one additional assumption on H , we show that it must have an internal (to H) non-branching path of the desired order.

Let ℓ denote the number of leaves in H which are not in path components and p denote the number of components in H which are paths. Suppose H is such (subject to our condition on $|\mathcal{C}|$ and that the number of components in H are minimized) that $\ell + p$ is minimized. Let L consist of the set of leaves in H , with one leaf selected from each path component. Note $|L| = \ell + p$. Note that there is a non-branching path of some order starting at each leaf. Let $\mathbf{v} = (v_1, v_2, \dots, v_n)$ be the vector where v_i is the number of leaves in L which are connected to a maximal non-branching path of order exactly i . Note, by assumption $v_i = 0$ for $i \geq 8$. We define the weight of \mathbf{v} to be:

$$wt(\mathbf{v}) = \sum_{i=1}^n v_i(1 - 2^{-i}).$$

Suppose that H is such that (subject to its other conditions) $wt(\mathbf{v})$ is minimized. We claim that $v_i \leq 2$ for all i . If $v_i \geq 3$, then applying the swapping lemma to the three vertices ending paths of order i would result in a new $H' = (H - \{x_i\}) \cup \{z\}$ and \mathcal{C}' where $wt(\mathbf{v}') < wt(\mathbf{v})$. If $\sum_{i \geq 4} v_i \geq 2$, we are done. If $\sum_{i \geq 4} v_i = 1$, then there are at most 7 leaves, and $\sum_i i v_i \leq 19$. Otherwise, there are at most 6 leaves and $\sum_i i v_i \leq 12$.

We also need the following lemma, whose proof we give after we complete the proof of Lemma 4.

Lemma 5. *Under the assumption that H does not contain two disjoint non-branching paths of order at least four, one of which starts at a leaf, we may assume that $|H| \geq 40$.*

Let J denote the set of vertices in H of degree at least 3 (in H), and let $j = |J|$. Let c denote the number of components of H . The vertices of J divide the $|H| - j$ vertices of $H - J$ into

$$c + \sum_{v \in J} (|N(v)| - 1) = c + j + \sum_{v \in J} (|N(v)| - 2) = j + \ell - c$$

segments which are non-branching paths, some of which may be empty if there are adjacent vertices in J .

Of these $j + \ell - c$ segments, $\ell + p$ of them are attached to vertices in L and these account for at most $\sum_i i v_i$ vertices. Thus, the remaining

$\geq |H| - 2 - j - \sum_i iv_i$ vertices are split into $j + \ell - c - (\ell + p)$ segments. If $\sum_{i \geq 4} v_i = 1$, so that $\ell \leq 7$ and $\sum_i iv_i \leq 19$, then there must be a non-branching path of order at least:

$$\left\lceil \frac{|H| - 2 - j - 19}{j - c - p} \right\rceil \geq \left\lceil \frac{|H| - 2 - (\ell - 2) - 19}{(\ell - 2) - c - p} \right\rceil \geq \left\lceil \frac{40 - 26}{4} \right\rceil = 4$$

since $|H| \geq 40$. Thus there is a non-branching path of order 4 in the interior of the forest, and along with one adjacent to a leaf we have the two desired paths of order 4.

If $\sum_{i \geq 4} v_i = 0$, so that $\ell \leq 6$ and $\sum_i iv_i \leq 12$, we have a non-branching path of order at least:

$$\left\lceil \frac{|H| - 2 - j - 12}{j - c - p} \right\rceil \geq \left\lceil \frac{|H| - 2 - (\ell - 2) - 12}{(\ell - 2) - c - p} \right\rceil \geq \left\lceil \frac{40 - 18}{3} \right\rceil = 8,$$

as desired. \square

Before we prove Lemma 5 we need one additional claim concerning cycles with edges between them. We define the graphs $G(a, b, c, d)$ to be the disjoint union of two cycles C_1 and C_2 of order a and b respectively, with two vertices v_1 and v_2 at distance d lying on cycle C_c specified. The vertex v_1 is connected to all of the other cycle, while v_2 is connected to a single vertex of the other cycle. (Note that for all choices of v_1, v_2 and the neighbor of v_2 , the resulting graphs are isomorphic).

Lemma 6. *Suppose C_1 and C_2 are two cycles with $|C_1| + |C_2| \geq 7$, and $e(C_1, C_2)$ edges between them. If $|e(C_1, C_2)| \geq 9$, then either $C_1 \cup C_2 \cup e(C_1, C_2) \subseteq G(|C_1|, |C_2|, c, d)$ for some parameters c and d , or there exist two shorter disjoint cycles.*

Note that 9 here is sharp: there exist configurations of C_1 and C_2 with 8 edges between them that admit no two shorter disjoint cycles, and are not contained in a $G(|C_1|, |C_2|, c, d)$.

Proof. Suppose $|e(C_1, C_2)| = 9$, and let v be a vertex with the largest number of edges into the other cycle. Without loss of generality, $v \in C_1$ and we let d_v denote the number of vertices in C_2 incident to v . We show $d_v \geq 8$. Note that this proves the lemma: if $|e(C_1, C_2)| > 9$, we consider groups of 9 edges at a time and it is clear that the vertex v of highest degree must be the same for all groups of edges.

We fix some order on C_2 . For vertices, x, y in C_2 , with $x \leq y$ (with respect to the order) we let (x, y) and $[x, y]$ denote the open or closed intervals of vertices from x to y . If $y \leq x$, then $[x, y]$ is the complement of (y, x) and similarly for (x, y) . Half open intervals, e.g. $(x, y]$, are defined accordingly.

Suppose $|e(v, C_2)| = 7$. Let $x_1 < x_2 < \dots < x_7 \in C_2$ denote the neighbors of v in order around the cycle. Then there are two other edges from C_1 to C_2 , which hit vertices $a_1 \leq a_2$ in C_2 (where a_1 may be the same as a_2). Either there are three x_i within $[a_1, a_2]$ or there are three x_i on the outside of $[a_1, a_2]$. In either case, one can build two shorter cycles which avoids (at least) one of the three vertices to the inside (or outside) of $[a_1, a_2]$.

If $|e(v, C_2)| = 6$, let $x_1 < \dots < x_6$ denote the vertices adjacent to v . Note that if $C_1 \setminus v$ has 2 neighbors in (x_2, x_6) then shorter cycles can be constructed avoiding x_2 . Otherwise $C_1 \setminus v$ has 2 neighbors in $[x_6, x_2]$ and shorter cycles can be constructed avoiding x_3 .

If $|e(v, C_2)| = 5$, let $x_1 < \dots < x_5$ denote the vertices adjacent to v . If $[x_5, x_1)$ contains 2 neighbors of $C_1 \setminus v$ then shorter cycles can be constructed avoiding x_2 . Similarly if there are two vertices in any of the intervals $[x_1, x_3)$, or $[x_3, x_5)$. But one of these must contain two elements.

If $|e(v, C_2)| = 4$, let $x_1 < \dots < x_4$ denote the vertices adjacent to v . Either one of $[x_4, x_1)$, $[x_1, x_2)$, $[x_2, x_3)$, or $[x_3, x_4)$ contains two edges from $C_1 \setminus v$, allowing a shorter cycle in any case or each has exactly one edge from $C_1 \setminus v$, which also allows a shorter cycle.

If $|e(v, C_2)| = 3$, let $x_1 < \dots < x_3$ denote the vertices adjacent to v . If one of (x_1, x_2) , (x_2, x_3) or (x_3, x_1) contains two edges from $C_1 \setminus v$, then G admits a shorter disjoint cycle. Now either one of x_1, x_2 , or x_3 has two edges from $C_1 \setminus v$ or there is exactly one edge from $C_1 \setminus v$ into each of x_1, x_2, x_3 , (x_1, x_2) , (x_2, x_3) , and (x_3, x_1) . In the second case, each set must be non-empty, and it is easy to see that there are shorter disjoint cycles. If x_1 (say) has two edges from $C_1 \setminus v$, then there are shorter disjoint cycles unless (x_3, x_1) and (x_1, x_2) are empty. This forces x_2 (say) to have two edges from $C_1 \setminus v$, forcing (x_2, x_3) to be empty, and thus x_3 must also have two edges from $C_1 \setminus v$. (Note that no x_i may have 3 edges from $C_1 \setminus v$, this would contradict the fact that v has maximum degree to the other cycle.) The same argument applied in reverse (now taking one of the x_i 's to be the vertex of degree 3, and looking at edges into C_2) implies that either we have two shorter cycles or, in fact, we must two triangles with a $K_{3,3}$ connecting them. This contradicts $|C_1| + |C_2| \geq 7$.

If $|e(v, C_2)| = 2$, let $x_1 < x_2$ denote the neighbors of v . If (x_1, x_2) or (x_2, x_1) have three edges from $C_1 \setminus v$ then there are shorter disjoint cycles. Thus one of x_1 or x_2 has at least 2 neighbors from $C_1 \setminus v$. This contradicts the fact that v has the maximum degree.

If $|e(v, C_2)| = 1$, suppose $v_1, v_2 \in C_1$ are as close together as possible such that v_1, v_2 have neighbors in C_2 , and let $x_1 < x_2$ denote their neighbors. One of (x_1, x_2) or (x_2, x_1) must have at least 3 neighbors in $C_1 \setminus \{v_1, v_2\}$ and thus there are shorter cycles.

Therefore $|e(v, C_2)| \geq 8$ as desired. □

Proof of Lemma 5. Suppose $|H| < 40$. Then the average size of a cycle in \mathcal{C} is at least $\frac{30k-40}{k-1} \geq 20$ (since $k > 1$). Let C be a cycle in \mathcal{C} of maximum order. Then C has no chords, due to the minimality of $|\mathcal{C}|$. Order the vertices in C , and denote them v_0, \dots, v_m . Let $R = \bar{C} = G - C$. We may assume that, without loss of generality at most two (adjacent) vertices in C have $|N_R(v)| < k - 1$, and if these exist, we will assume that they are (possibly only one of) v_1, v_2 . Pair off vertices $\{\{v_{4i}, v_{4i+2}\}, \{v_{4i+1}, v_{4i+3}\} : i = 0, 1, \dots, \lfloor m/4 \rfloor - 1\}$ as long as it is possible to form both pairs. Let U denote the set of vertices not paired off; there are $m + 1 \pmod{4}$ such vertices. Then for any of these pairs (u, v) , $|N_C(u, v)| = 3$. Now consider that:

$$\sum_{i \leq \lfloor m/4 \rfloor - 1} (|N_R(v_{4i}, v_{4i+2})| + |N_R(v_{4i+1}, v_{4i+3})|) + \sum_{v \in U} |N_R(v)| \geq (m+1)(k-1).$$

Note that $|e(C, H)| < 20$ since if any two vertices at distance at most 9 in H have neighbors in C , then there is a shorter cycle. Here we use the fact that H is minimized with respect to $wt(\mathbf{v})$ and the condition on the non-branching paths within H . Note that this is already a contradiction if $k = 2$. Thus there exists some cycle $C' \in \mathcal{C}$, with at least $m + 2$ edges between C and C' .

Lemma 6, along with the neighborhood condition, implies that $|C'| = |C|$ and that a vertex in C must be adjacent to every vertex in C' , with one additional edge between the two. The fact that the high degree vertex is in C comes from the neighborhood condition, and the fact that $|C'| = |C|$ comes from the fact that there is a vertex in C with degree $m + 1$ into C' , and C is a maximum order cycle.

We now repeat the argument: Let v be the vertex in C of high degree

to C' . We pair off vertices of C starting at the successor of v , and also pair off the vertices of C' to get that there must be at least $(2m + 1)(k - 1) - 1$ edges leaving $C \cup C' \setminus v$; in particular there must be third cycle C'' such that there are at least $2m + 1$ edges between C and C' and C'' . Applying Lemma 6 again implies $|C''| = |C'| = |C|$ and that there is a $v' \neq v \in C$ and a $v'' \in C'$ each of which are adjacent to at least $|C''| - 1$ vertices in $|C''|$. But then it is clear that we can find three shorter cycles, contradicting our assumptions on $|C|$. (Here, if $k = 3$ the number of edges between $C \cup C'$ and H already contradict the fact that $|C|$ is minimized.) \square

We are now ready to complete the proof of Theorem 2. By Lemma 4, we may assume that H contains either two disjoint non-branching paths of order 4, or one non-branching path of order 8. First consider the case where H contains two disjoint non-branching paths P_1, P_2 of order 4 (one of which contains a leaf). Let $P_1 = \{x_1, x_2, x_3, x_4\}$ and $P_2 = \{y_1, y_2, y_3, y_4\}$, and consider the pairs of non-adjacent vertices $S = \{(x_1, x_3), (x_2, x_4), (y_1, y_3), (y_2, y_4)\}$. Then

$$\sum_{(u,v) \in S} |N_C(u, v)| \geq 4 \cdot (2k - 2) + 1 = 8(k - 1) + 1.$$

Therefore, there exists a $C = C_i$ such that $\sum_{(u,v) \in S} |N_C(u, v)| \geq 9$. This implies, by Claim 3 that some vertex in one of the P_i s has two neighbors on C , so $|C| \leq 4$.

First, assume that $|C| = 4$. Without loss of generality, assume that

$$|N_C(x_1, x_3)| + |N_C(x_2, x_4)| \geq |N_C(y_1, y_3)| + |N_C(y_2, y_4)|.$$

If $|N_C(x_1, x_3)| + |N_C(x_2, x_4)| \geq 7$, then one pair, say x_1 and x_3 is adjacent to all four vertices on C . Then we reach a contradiction to the minimization of $|C|$ as either a triangle is already formed, or x_2 or x_4 must be adjacent to some vertex on the cycle, creating a triangle. If $|N_C(x_1, x_3)| + |N_C(x_2, x_4)| = 6$, then there are two vertices on C which are adjacent to two vertices in P_1 , and these vertices must come from different classes, i.e. one from $\{x_1, x_3\}$ and one from $\{x_2, x_4\}$, and either both other vertices in C are adjacent to one other vertex on P_1 , or one vertex is adjacent to two vertices on P_1 . Suppose first two vertices on C are adjacent to two vertices of P_1 , and the others are adjacent to one vertex. Then the two vertices of C which are adjacent to two vertices of P_1 must be adjacent to the two end vertices of P_1 (to avoid a triangle since they must come from different classes) and must be antipodal. Label the vertices of C to be (in order) c_1, \dots, c_4 . Then without

loss of generality x_1 and x_4 are adjacent to c_1 and c_3 . We are done unless both c_2 and c_4 are adjacent to one of x_2 or x_3 . In this case, we must use P_2 . We know that $|N_C(y_1, y_3)| + |N_C(y_2, y_4)| \geq 3$. Therefore either some vertex in C is adjacent to two vertices of P_2 (in which case it is easy to find disjoint cycles), or three different vertices of C are adjacent to P_2 . It is easy to see here that however they are adjacent we are done, though we note that 2 edges does not suffice here: if P_2 were just adjacent to c_1 and c_3 we could not find the shorter disjoint cycles. This is the only one of these cases where we use the ninth edge, and hence requires care when we consider the case where we have a single path of length 8 below.

If $|N_C(x_1, x_3)| + |N_C(x_2, x_4)| = 5$, then some vertex z in C is adjacent to two vertices of P_1 , and these must be the end vertices. If P_2 is adjacent to two vertices that are not z , then we have two disjoint cycles. If this doesn't occur, z occurs in the neighborhood of 2 vertices in P_2 (again, the end vertices) as in total P_2 sends three edges to C . But then P_1 has two edges that don't go to z , leading to two disjoint cycles (one involving P_1 and vertices on $C - \{z\}$ and one involving P_2 and $\{z\}$). We also will handle the case where $|N_C(x_1, x_3)| + |N_C(x_2, x_4)| = 4$, which arises in the case where there is a non-branching path of order 8. If $|N_C(x_1, x_3)| + |N_C(x_2, x_4)| = 4$, then either some vertex is adjacent to two vertices of P_1 or two vertices of P_2 which leads to the exact case above, or all vertices are adjacent to exactly one vertex in both P_1 and P_2 . In either case, there are two disjoint cycles and we are done.

If $|C| = 3$, then note $|N_C(x_1, x_3)| + |N_C(x_2, x_4)| \leq 6$ (again, we assume that P_1 has the larger of the neighborhoods). If this is at least 5, two vertices on C must be adjacent to two vertices of P_1 , and the other vertex on C must be adjacent to (at least) one vertex on P_1 . If P_2 sends three edges into C , we are clearly done. Either two hit some vertex, and we have a cycle involving P_1 and the vertices not hit by P_2 , or all three are hit and we have an analogous situation using one of the vertices that P_2 hits twice. If P_2 only sends 2 edges into C , then all three of the vertices of C must be hit twice by P_1 and, again, we have two disjoint cycles. If $|N_C(x_1, x_3)| + |N_C(x_2, x_4)| = 4$, then some vertex z of C must be adjacent to two vertices in P_1 . Two of the edges from P_2 must be incident to vertices different than z , and thus we have two disjoint cycles as desired. Again, this last case cannot occur unless we are in the non-branching path of order 8 case.

Thus we may assume we are in the case where there is a single non-branching path of order 8; let $P = \{x_1, \dots, x_8\}$. We can think of $P =$

$P_1 \cup P_2$, where P_1 and P_2 are the first and last four vertices of P respectively. The same argument as above guarantees that there is a C such that $\sum_{(u,v) \in S} |N_C(u,v)| \geq 8$. This is where we lose something: having a leaf in our path gave us one extra edge. Note that before we used that extra edge to claim that $|C| \leq 4$, but did not use the extra edge to produce the two disjoint cycles, we showed their existence even when the (sum of the) neighborhoods of both P_1 and P_2 was of size 4. Note either every vertex has $|N_C(x_i)| = 1$ with $N_C(x_i) \neq N_C(x_{i+2})$, or $|C| \leq 4$. If $|C| \leq 4$, then we can apply the above arguments, noting that we have already handled the additional cases which occur except for one noted.

The last case with $|C| \leq 4$ is when $|C| = 4$, and we have $C = \{c_1, c_2, c_3, c_4\}$ and $P_1 = \{x_1, x_2, x_3, x_4\}$ where x_1 and x_4 are adjacent to both c_1 and c_3 with either x_2 or x_3 adjacent to both c_2 and c_4 . Then there must be at least two edges between P_2 and C ; if there are three then the argument above shows there exist two disjoint cycles. The only case that the above argument does not cover is if the neighbors of P_2 are c_1 and c_3 . But even in this case we are done as P_1 and P_2 are adjacent: we take the cycle involving x_1, c_1, c_2 , the neighbor of c_2 on P_1 (either x_2 or x_3) and the segment of P_1 connecting this vertex and x_1 and also the cycle involving x_4, c_3 , the neighbor of c_3 in P_2 and the path between x_4 and this vertex.

If every vertex has degree 1, consider that x_1 and x_3 have two distinct neighbors on C . If $N_C(x_2) \in \{N_C(x_1), N_C(x_3)\}$, then we have a triangle, and hence $|C| = 3 \leq 4$ and we are already done. Now assume $N_C(x_2) \notin \{N_C(x_1), N_C(x_3)\}$. Order the vertices along the cycle so that $N_C(x_1) < N_C(x_2) < N_C(x_3)$. Then if $N_C(x_4) \notin [N_C(x_1), N_C(x_2)]$ we are done, so we may assume that $N_C(x_4)$ is in this interval.

Now consider the neighbors of x_i for $i \geq 5$. If $N_C(x_i) \notin [N_C(x_1), N_C(x_2)]$ we are done as above. In fact, if $N_C(x_i) \in (N_C(x_1), N_C(x_2))$ we are also done: we may use x_4, x_i and the portions on the path and cycle connecting them for one cycle, using the portion of the cycle contained in $[N_C(x_1), N_C(x_2))$. Then one of $N_C(x_1)$ or $N_C(x_2)$ will not be used in that cycle; we will use that vertex, along with $N_C(x_3)$ the portion connecting them in the cycle not containing $(N_C(x_1), N_C(x_2))$ along with their neighbors on the path and the portion of the path connecting them. Thus all vertices x_i for $i \geq 5$ must have their neighbors be either x_1 or x_2 . But then one of these vertices will have two neighbors x_i with $i \geq 5$. This easily admits two disjoint cycles.

□

References

- [1] K. Corrádi, A. Hajnal, On the maximal number of independent circuits in a graph, *Acta Math. Acad. Sci. Hungar.* **14** (1963), 423–439.
- [2] J. Faudree, R. J. Gould, A note on neighborhood unions and independent cycles, *Ars Combinatoria*, **76** (2005), 29–31.
- [3] D. Finkel, On the number of independent chorded cycles in a graph, *Discrete Math.*, **308** (2008), Issue 22, 5265–5268.
- [4] P. Justesen, On independent circuits in finite graphs and a conjecture of Erdős and Posa, *Annals of Discrete Math.* **41** (1989), 299–305.
- [5] H. Wang, On the maximum number of independent cycles in a graph, *Discrete Math.* **205** (1999), Issues 1-3, 183–190.